


复几何课程

Lecture 1. Holomorphic functions of several complex variables.

$$\bar{\partial} f = 0$$

$$f(z_1, \dots, z_n) = \sum c_\nu z^\nu$$



Outline (2 hours)

1. ∂ & $\bar{\partial}$

2. Cauchy integral formula $\left\{ \begin{array}{l} \text{dim 1} \\ \text{higher dim} \end{array} \right.$

3. Holomorphic functions and their properties

4. Hartogs Theorem (Prove by solving $\bar{\partial}$ -equ)

Power series expansion
Weierstrass's convergence thm
Cauchy's ineq.
Max principle
Identity theorem

• 1. ∂ & $\bar{\partial}$

Let $\Omega \subseteq \mathbb{C}^n$ be a domain (open & connected subset in \mathbb{C}^n)

Let (z^1, \dots, z^n) be the standard complex coordinates of \mathbb{C}^n .

Put $z^i = x^i + \sqrt{-1} y^i$.

Define ∂ & $\bar{\partial}$ as follows.

$$\begin{cases} \frac{\partial}{\partial z^i} := \frac{1}{2} \left(\frac{\partial}{\partial x^i} - \sqrt{-1} \frac{\partial}{\partial y^i} \right) \\ \frac{\partial}{\partial \bar{z}^i} := \frac{1}{2} \left(\frac{\partial}{\partial x^i} + \sqrt{-1} \frac{\partial}{\partial y^i} \right) \end{cases} \quad \& \quad \begin{cases} dz^i = dx^i + \sqrt{-1} dy^i \\ d\bar{z}^i = dx^i - \sqrt{-1} dy^i \end{cases} \quad \text{for } \forall 1 \leq i \leq n.$$

$$\text{Then } \begin{cases} \partial := \sum_{i=1}^n \frac{\partial}{\partial z^i} \otimes dz^i \\ \bar{\partial} := \sum_{i=1}^n \frac{\partial}{\partial \bar{z}^i} \otimes d\bar{z}^i \end{cases}$$

e.g. Check that $\frac{\sqrt{-1}}{2} dz^i \wedge d\bar{z}^i = dx^i \wedge dy^i$.

More concretely, for $\forall f \in C^1(\Omega, \mathbb{C})$, we have

$$\begin{cases} \partial f = \sum_i \frac{\partial f}{\partial z^i} dz^i = \frac{1}{2} \sum_i \left(\frac{\partial f}{\partial x^i} dx^i + \frac{\partial f}{\partial y^i} dy^i + \sqrt{-1} \frac{\partial f}{\partial x^i} dy^i - \sqrt{-1} \frac{\partial f}{\partial y^i} dx^i \right) \\ \bar{\partial} f = \sum_i \frac{\partial f}{\partial \bar{z}^i} d\bar{z}^i = \frac{1}{2} \sum_i \left(\frac{\partial f}{\partial x^i} dx^i + \frac{\partial f}{\partial y^i} dy^i + \sqrt{-1} \frac{\partial f}{\partial y^i} dx^i - \sqrt{-1} \frac{\partial f}{\partial x^i} dy^i \right) \end{cases}$$

▲ Observe that

$$\partial f + \bar{\partial} f = df = \sum_i \left(\frac{\partial f}{\partial x^i} dx^i + \frac{\partial f}{\partial y^i} dy^i \right).$$

Namely, $d = \partial + \bar{\partial}$

★ ▲ We call $f \in C^1(\Omega, \mathbb{C}^n)$ holomorphic if f satisfies

$$\bar{\partial} f = 0, \text{ ie if we write } f = u + \sqrt{-1}v \text{ then}$$

$$\sum_i \left(\frac{\partial u}{\partial x^i} dx^i + \sqrt{-1} \frac{\partial v}{\partial x^i} dx^i + \frac{\partial u}{\partial y^i} dy^i + \sqrt{-1} \frac{\partial u}{\partial y^i} dy^i + \sqrt{-1} \frac{\partial u}{\partial y^i} dx^i - \frac{\partial v}{\partial y^i} dx^i - \sqrt{-1} \frac{\partial u}{\partial x^i} dy^i + \frac{\partial v}{\partial x^i} dy^i \right) = 0$$

$$\Leftrightarrow \begin{cases} \frac{\partial u}{\partial x^i} = \frac{\partial v}{\partial y^i} \\ \frac{\partial u}{\partial y^i} = -\frac{\partial v}{\partial x^i} \end{cases} \text{ for } \forall 1 \leq i \leq n.$$

This is called Cauchy-Riemann equation.

In particular, if f is holomorphic, then f is holomorphic in each complex variable (as a one-variable holomorphic/analytic function).

• 2 Cauchy integral formula.

- When $n=1$, assume that Ω is a bounded open set in \mathbb{C} s.t. $\partial\Omega$ consists of finitely many C^1 Jordan curves. Then for $\forall u \in C^1(\bar{\Omega})$ we have

$$u(z_0) = \frac{1}{2\pi i} \left(\int_{\partial\Omega} \frac{u(z)}{z-z_0} dz + \iint_{\Omega} \frac{\partial u / \partial \bar{z}}{z-z_0} dz \wedge d\bar{z} \right) \text{ for } \forall z_0 \in \Omega.$$

pf: By definition

$$\begin{aligned} \iint_{\Omega} \frac{\frac{\partial u}{\partial \bar{z}} dz \wedge d\bar{z}}{(z-z_0)} &= \lim_{\epsilon \rightarrow 0} \iint_{\Omega \setminus B_{\epsilon}(z_0)} \frac{\frac{\partial u}{\partial \bar{z}} dz \wedge d\bar{z}}{(z-z_0)} \\ &= \lim_{\epsilon \rightarrow 0} \iint_{\Omega \setminus B_{\epsilon}(z_0)} \frac{-\bar{\partial}(u dz)}{(z-z_0)} = \lim_{\epsilon \rightarrow 0} \iint_{\Omega \setminus B_{\epsilon}(z_0)} \frac{-d(u dz)}{(z-z_0)} \\ &= \lim_{\epsilon \rightarrow 0} \left(\int_{\partial B_{\epsilon}(z_0)} \frac{u dz}{z-z_0} - \int_{\partial\Omega} \frac{u dz}{z-z_0} \right) \\ &= \lim_{\epsilon \rightarrow 0} \int_{\partial B_{\epsilon}(z_0)} \frac{u(z_0) dz}{z-z_0} - \int_{\partial\Omega} \frac{u dz}{z-z_0} \\ &= 2\pi i u(z_0) - \int_{\partial\Omega} \frac{u dz}{z-z_0} \end{aligned}$$

This completes the proof. \square

▲ The above formula has several consequences.

$n=1$ { ① When f is holomorphic ($n=1$) one has

$$f(z_0) = \frac{1}{2\pi i} \int_{\partial\Omega} \frac{f(z)}{z-z_0} dz$$
, which is the classical Cauchy integral formula.

② When $f \in C^1(\Omega)$, then

$$f(z_0) = \frac{1}{2\pi i} \iint_{\Omega} \frac{\partial f / \partial \bar{z}}{z-z_0} dz \wedge d\bar{z}$$

$n \geq 1$ — ③ For general $n \geq 1$, if f is holomorphic on $\{(z_1, \dots, z_n) \in \mathbb{C}^n \mid |z_i| < r_i, i=1, \dots, n\} =: \mathbb{D}_r$
 $r = (r_1, \dots, r_n) \in (\mathbb{R}_{>0})^n$
 then $f(\xi) = \left(\frac{1}{2\pi i}\right)^n \int_{|z^1|=r_1} \dots \int_{|z^n|=r_n} \frac{f(z^1, \dots, z^n)}{(z^1-\xi^1) \dots (z^n-\xi^n)} dz^1 \dots dz^n$ for $\forall \xi = (\xi^1, \dots, \xi^n) \in \mathbb{D}_r$

This is the Cauchy integral formula for holomorphic functions of several cplx varb.

• Def: We put $\mathcal{O}(\Omega) := \{ \text{holomorphic function on } \Omega \}$.

• Thm. The following are equivalent

① $f \in \mathcal{O}(\Omega)$

② f satisfies the Cauchy integral formula for \forall polydisk $D_r \subset \Omega$.

③ For $\forall z_0 \in \Omega$, \exists polydisk D_r around z_0 s.t.

$$f(z) = \sum_{\nu \in \mathbb{N}^n} a_\nu (z - z_0)^\nu \quad \text{namely } f \text{ has a power series expansion}$$

• Prnk In ③, a_ν is given by

$$a_\nu = \left(\frac{1}{2\pi i}\right)^n \int_{|z^1 - z_0^1| = r_1} \dots \int_{|z^n - z_0^n| = r_n} \frac{f(z^1, \dots, z^n)}{(z^1 - z_0^1)^{\nu_1+1} \dots (z^n - z_0^n)^{\nu_n+1}} dz^1 \dots dz^n$$

which is indep. of the choice of polydisk as long as $|r| \ll 1$.

$$a_\nu = \frac{1}{\nu_1! \dots \nu_n!} \frac{\partial^{|\nu|} f}{\partial z_1^{\nu_1} \dots \partial z_n^{\nu_n}}(z_0) =: D^\nu f(z_0).$$

This follows from the Taylor expansion

$$\frac{1}{(z - z_0)} = \frac{1}{z - z_0 - (z - z_0)} = \frac{1}{(z - z_0)} \frac{1}{1 - \frac{z - z_0}{z - z_0}} = \frac{1}{(z - z_0)} \sum_{\nu} \left(\frac{z - z_0}{z - z_0}\right)^\nu$$

two convergent power series around $0 \in \mathbb{C}^n$

& if $f = g$ over some small abhd $V \subset \mathbb{C}^n$ containing 0

then $a_\nu = b_\nu$.

(Since a_ν & b_ν are determined by the infinitesimal information of $f+g$ around 0)

• Thm Weierstrass's Convergence theorem.

Let $\{f_k\} \subset \mathcal{O}(\Omega)$ be a sequence of hol. functions on Ω that converges uniformly to a function f . Then $f \in \mathcal{O}(\Omega)$.

pf: $f = \lim f_k = \lim \int \int \frac{f_k}{(z-\zeta)} d\zeta = \int \int \frac{f}{(z-\zeta)} d\zeta$. \square

• Thm. For $f_1, f_2 \in \mathcal{O}(\Omega)$, assume that for some $U \subseteq \Omega$
 $f_1|_U = f_2|_U$. then $f_1 = f_2$.

pf: Put $N := \{z_0 \in \Omega \text{ s.t. } D^v f_1(z_0) = D^v f_2(z_0) \text{ for } \forall v \in \mathbb{N}^n\}$.

Then N is clearly closed. & $U \subseteq N$.

N is also open as $D^v f_1(z_0) = D^v f_2(z_0)$ implies that $f_1 = f_2$ around z_0 .
Thus we must have $N = \Omega$.

• Thm (Max principle). If $f \in \mathcal{O}(\Omega)$ & $\exists z_0 \in \Omega$ s.t.

$|f|$ is locally maximized at z_0 , then f is constant. \square

pf 1. Consider complex lines through z_0 & use max principle of 1-variable hol. funct.

pf 2. Using mean value formula of f & using the fact that $|f| = \text{const} \Rightarrow f = \text{const}$.

• Thm (Hartogs thm) Assume that $n \geq 2$.

Let Ω be a domain & $K \subseteq \Omega$ a compact subset s.t. $\Omega \setminus K$ connected.

Then $\forall f \in \mathcal{O}(\Omega \setminus K)$ can be extended to a function $\hat{f} \in \mathcal{O}(\Omega)$
 s.t. $f = \hat{f}$ on $\Omega \setminus K$.

▲ This is obviously not true when $n=1$.

pf: Choose a cut-off function $\varphi \in C_0^\infty(\Omega)$ s.t.
 $\varphi \equiv 1$ on a nbhd of K . Consider

$f_0 := (1-\varphi)f$. Then $f_0 \in C^\infty(\Omega)$

Put $\alpha := \bar{\partial} f_0 = -f \bar{\partial} \varphi$, which is a C_0^∞ - $(0,1)$ form on \mathbb{C}^n .

Obviously, $\bar{\partial} \alpha = 0$. If write $\alpha = \sum_{i=1}^n \alpha_i d\bar{z}_i$

then $\frac{\partial \alpha_i}{\partial \bar{z}_j} = \frac{\partial \alpha_j}{\partial \bar{z}_i}$ for $\forall i=j$.

Put $u(z) = \frac{1}{2\pi i} \int_{B_R} \frac{\alpha_i(\tau, z_2, \dots, z_n)}{(\tau - z_1)} d\tau \wedge d\bar{\tau}$ for $\forall z \in \Omega$.

Fix a ball $B_R \subseteq \mathbb{C}$ with R sufficiently large. \leftarrow
 $= \frac{1}{2\pi i} \int_{B_R} \frac{\alpha_i(\tau + z_1, z_2, \dots, z_n)}{\tau} d\tau \wedge d\bar{\tau}$

So in particular $u \in C_0^\infty$ & we have

$\frac{\partial u(z)}{\partial \bar{z}_k} = \frac{1}{2\pi i} \int_{B_R} \frac{\partial \alpha_k(\tau, z_2, \dots, z_n) / \partial \bar{z}_k}{(\tau - z_1)} d\tau \wedge d\bar{\tau} = \alpha_k(z)$ by Cauchy integral formula.

Thus u solves $\bar{\partial}u = \alpha \Rightarrow \bar{\partial}(f_0 - u) = 0$ on Ω .

$$\Rightarrow \hat{f} := f_0 - u \in \mathcal{O}(\Omega)$$

Notice that $\begin{cases} u=0 \\ f_0=f \end{cases}$ on an open subset of $\Omega \setminus K$, thus $\hat{f} = f$ on an open subset of $\Omega \setminus K$ & hence $\hat{f} = f$ on $\Omega \setminus K$ (as $\Omega \setminus K$ is connected)

This completes the proof. \square .

