

Motivic decompositions for the Hilbert scheme of points of a K3 surface

By *Andrei Neguț* at Cambridge, MA, *Georg Oberdieck* at Bonn and *Qizheng Yin* at Beijing

Abstract. We construct an explicit, multiplicative Chow–Künneth decomposition for the Hilbert scheme of points of a K3 surface. We further refine this decomposition with respect to the action of the Looijenga–Lunts–Verbitsky Lie algebra.

1. Introduction

In the present paper, we study the motivic aspects of the Looijenga–Lunts–Verbitsky ([20, 34], LLV for short) Lie algebra action on the Chow ring of the Hilbert scheme of points of a K3 surface. Using a special element of the LLV algebra and formulas of [23] by Maulik and the first author, we construct an explicit Chow–Künneth decomposition for the Hilbert scheme, prove its multiplicativity, and show that all divisor classes and Chern classes lie in the correct component of the decomposition. This confirms expectations of Beauville [2] and Voisin [36]. We also obtain a refined motivic decomposition for the Hilbert scheme by taking into account the LLV algebra action, and prove its multiplicativity.

Both results parallel the case of an abelian variety, which we shall briefly review.

1.1. Abelian varieties. Let X be an abelian variety of dimension g . Recall the classical result of Deninger–Murre on the decomposition of the Chow motive $\mathfrak{h}(X)$.

Theorem 1.1 ([5]). *There is a unique, multiplicative Chow–Künneth decomposition*

$$(1.1) \quad \mathfrak{h}(X) = \bigoplus_{i=0}^{2g} \mathfrak{h}^i(X)$$

such that for all $N \in \mathbb{Z}$, the multiplication $[N] : X \rightarrow X$ acts on $\mathfrak{h}^i(X)$ by $[N]^ = N^i$.*

The corresponding author is Andrei Neguț.

Andrei Neguț gratefully acknowledges the NSF grants DMS-1760264 and DMS-1845034, as well as support from the Alfred P. Sloan Foundation. Qizheng Yin was supported by the NSFC grants 11701014, 11831013, and 11890661.

The decomposition in (1.1) specializes to the Künneth decomposition in cohomology (hence the name Chow–Künneth), and to the Beauville decomposition [1] in Chow. The latter takes the form

$$(1.2) \quad A^*(X) = \bigoplus_{i,s} A^i(X)_s$$

with

$$A^i(X)_s = A^i(\mathfrak{h}^{2i-s}(X)) = \{\alpha \in A^i(X) \mid [N]^*\alpha = N^{2i-s}\alpha \text{ for all } N \in \mathbb{Z}\}.$$

The *multiplicativity* of (1.1) stands for the fact that the cup product

$$\cup : \mathfrak{h}(X) \otimes \mathfrak{h}(X) \rightarrow \mathfrak{h}(X)$$

respects the grading, in the sense that

$$\cup : \mathfrak{h}^i(X) \otimes \mathfrak{h}^j(X) \rightarrow \mathfrak{h}^{i+j}(X) \quad \text{for all } i, j \in \{0, \dots, 2g\}.$$

This can be seen by simply comparing the actions of $[N]^*$. As a result, the bigrading in (1.2) is multiplicative, i.e., compatible with the ring structure of $A^*(X)$.

The Beauville decomposition is expected to provide a multiplicative *splitting* of the conjectural Bloch–Beilinson filtration on $A^*(X)$. A difficult conjecture of Beauville (and consequence of the Bloch–Beilinson conjecture) predicts the vanishing $A^*(X)_s = 0$ for $s < 0$ and the injectivity of the cycle class map

$$\text{cl} : A^*(X)_0 \rightarrow H^*(X).$$

Further, any symmetric ample class $\alpha \in A^1(X)_0$ induces an \mathfrak{sl}_2 -triple (e_α, f_α, h) acting on $A^*(X)$. A Lefschetz decomposition of $\mathfrak{h}(X)$ with respect to the \mathfrak{sl}_2 -action was obtained by Künnemann [16], refining (1.1). More generally, Moonen [24] constructed an action of the Néron–Severi part of the Looijenga–Lunts [20] Lie algebra \mathfrak{g}_{NS} on $A^*(X)$, which contains all possible \mathfrak{sl}_2 -triples above (he actually considered the slightly larger Lie algebra $\mathfrak{sp}(X \times X^\vee)$; see [24, Section 6]). He then obtained a refined motivic decomposition with respect to the \mathfrak{g}_{NS} -action.

Theorem 1.2 ([24]). *There is a unique decomposition*

$$(1.3) \quad \mathfrak{h}(X) = \bigoplus_{\psi \in \text{Irrep}(\mathfrak{g}_{\text{NS}})} \mathfrak{h}_\psi(X),$$

where ψ runs through all isomorphism classes of finite-dimensional irreducible representations of \mathfrak{g}_{NS} , and $\mathfrak{h}_\psi(X)$ is ψ -isotypic under \mathfrak{g}_{NS} .

Here being ψ -isotypic means that $\mathfrak{h}_\psi(X)$ is stable under \mathfrak{g}_{NS} and that for any Chow motive M , the \mathfrak{g}_{NS} -representation $\text{Hom}(M, \mathfrak{h}_\psi(X))$ is isomorphic to a direct sum of copies of ψ .

Again (1.3) specializes to refined decompositions in cohomology and in Chow.

1.2. Chow–Künneth. We switch to the case of the Hilbert scheme. Let S be a projective K3 surface over an algebraically closed field of characteristic 0, and let $X = \text{Hilb}_n(S)$ be the Hilbert scheme of n points on S .

In the paper [29], the second author lifted the action of the Néron–Severi part of the LLV algebra \mathfrak{g}_{NS} from cohomology to Chow. In particular, there is an explicit grading operator

$$h \in A^{2n}(X \times X)$$

which appears in every \mathfrak{sl}_2 -triple (e_α, f_α, h) in \mathfrak{g}_{NS} . We normalize h so that it acts on $H^{2i}(X)$ by multiplication by $i - n$.

We regard h as a natural replacement for the operator $[N]^*$ in the abelian variety case. Our first result decomposes the Chow motive $\mathfrak{h}(X)$ into eigenmotives of h .

Theorem 1.3. *There is a unique Chow–Künneth decomposition*

$$(1.4) \quad \mathfrak{h}(X) = \bigoplus_{i=0}^{2n} \mathfrak{h}^{2i}(X)$$

such that h acts on $\mathfrak{h}^{2i}(X)$ by multiplication by $i - n$.

The mutually orthogonal projectors in the decomposition (1.4) are written explicitly in terms of the Heisenberg algebra action [15, 26]. We also show that (1.4) agrees with the Chow–Künneth decomposition obtained by de Cataldo–Migliorini [4] and Vial [35].

As before the decomposition in (1.4) specializes to a decomposition in Chow:

$$(1.5) \quad A^*(X) = \bigoplus_{i,s} A^i(X)_{2s}$$

with

$$A^i(X)_{2s} = A^i(\mathfrak{h}^{2i-2s}(X)) = \{\alpha \in A^i(X) \mid h(\alpha) = (i - s - n)\alpha\}.$$

1.3. Multiplicativity. In the seminal paper [2], Beauville raised the question of whether hyper-Kähler varieties behave similarly to abelian varieties in the sense that the conjectural Bloch–Beilinson filtration also admits a multiplicative splitting. As a test case, he conjectured that for a hyper-Kähler variety, the cycle class map is injective on the subring generated by divisor classes.

For the Hilbert scheme of points of a K3 surface, Beauville’s conjecture was recently proven in [23]; see also [29] for a shorter proof. But the ultimate goal remains to find the multiplicative splitting. Meanwhile, Shen and Vial [32, 33] introduced the notion of a *multiplicative Chow–Künneth decomposition*, upgrading Beauville’s question from Chow groups to the level of correspondences/Chow motives.

The main result of this paper confirms that (1.4) provides a multiplicative Chow–Künneth decomposition for the Hilbert scheme.

Theorem 1.4. *Let S be a projective K3 surface and let $X = \text{Hilb}_n(S)$.*

(i) *The Chow–Künneth decomposition (1.4) is multiplicative, i.e., the cup product*

$$\cup : \mathfrak{h}(X) \otimes \mathfrak{h}(X) \rightarrow \mathfrak{h}(X)$$

respects the grading, in the sense that

$$\cup : \mathfrak{h}^{2i}(X) \otimes \mathfrak{h}^{2j}(X) \rightarrow \mathfrak{h}^{2i+2j}(X)$$

for all $i, j \in \{0, \dots, 2n\}$. As a result, the bigrading in (1.5) is multiplicative.

(ii) *All divisor classes and Chern classes of X belong to $A^*(X)_0$.*

Part (ii) of Theorem 1.4 is related to the Beauville–Voisin conjecture [36], which predicts that for a hyper-Kähler variety, the cycle class map is injective on the subring generated by divisor classes and Chern classes. In the Hilbert scheme case, one may further ask the vanishing $A^*(X)_{2s} = 0$ for $s < 0$ and the injectivity of the cycle class map

$$\text{cl} : A^*(X)_0 \rightarrow H^*(X).$$

We do not tackle these questions in the present paper.

The key to the proof of Theorem 1.4 (i) is the compatibility between the grading operator h and the cup product. For example, at the level of Chow groups, we show that the operator

$$\tilde{h} = h + n\Delta_X \in A^{2n}(X \times X)$$

acts on $A^*(X)$ by derivations, i.e.,

$$(1.6) \quad \tilde{h}(x \cdot x') = \tilde{h}(x) \cdot x' + x \cdot \tilde{h}(x')$$

for all $x, x' \in A^*(X)$. We achieve this by explicit calculations using the Chow lifts [23] of the well-known machinery for the Heisenberg algebra action [18, 19], and our argument yields (1.6) at the level of correspondences; see Section 4. Once the compatibility is established, part (i) of Theorem 1.4 is deduced by simply comparing the eigenvalues of \tilde{h} .

1.4. Previous work. Theorem 1.4 was previously obtained by Vial in [35] based on Voisin’s announced result [37, Theorem 5.12] on *universally defined cycles*. A second proof, also relying on Voisin’s theorem, was given by Fu and Tian [8]. They interpreted part (i) of Theorem 1.4 as the motivic incarnation of Ruan’s crepant resolution conjecture [31]. Our proof has the advantage of being explicit and unconditional at the moment.

We note that multiplicative Chow–Künneth decompositions, for both hyper-Kähler and non-hyper-Kähler varieties, have been studied in [6, 7, 9–12, 17].

1.5. Refined decomposition. We further obtain a refined decomposition of the Chow motive $\mathfrak{h}(X)$ with respect to the action of the Néron–Severi part of the LLV algebra \mathfrak{g}_{NS} . Both the statement and the proof parallel the abelian variety case.

Theorem 1.5. *Let S be a projective K3 surface and let $X = \text{Hilb}_n(S)$. There is a unique decomposition*

$$(1.7) \quad \mathfrak{h}(X) = \bigoplus_{\psi \in \text{Irrep}(\mathfrak{g}_{\text{NS}})} \mathfrak{h}_{\psi}(X),$$

where ψ runs through all isomorphism classes of finite-dimensional irreducible representations of \mathfrak{g}_{NS} , and $\mathfrak{h}_{\psi}(X)$ is ψ -isotypic under \mathfrak{g}_{NS} .

Consider the weight decomposition

$$\mathfrak{g}_{\text{NS}} = \mathfrak{g}_{\text{NS},-2} \oplus \mathfrak{g}_{\text{NS},0} \oplus \mathfrak{g}_{\text{NS},2}, \quad \mathfrak{g}_{\text{NS},0} = \bar{\mathfrak{g}}_{\text{NS}} \oplus \mathbb{Q} \cdot h,$$

where $\bar{\mathfrak{g}}_{\text{NS}}$ is the Néron–Severi part of the *reduced LLV algebra* (terminology taken from [14]). Let $\bar{\mathfrak{t}} \subset \bar{\mathfrak{g}}_{\text{NS}}$ be a Cartan subalgebra and write

$$\mathfrak{t} = \bar{\mathfrak{t}} \oplus \mathbb{Q} \cdot \tilde{h}.$$

Then the decomposition in (1.7) implies a motivic decomposition in terms of the irreducible representations (i.e., characters) of \mathfrak{t} as follows:

$$(1.8) \quad \mathfrak{h}(X) = \bigoplus_{\lambda \in \mathfrak{t}^*} \mathfrak{h}_\lambda(X).$$

The decomposition (1.8) specializes to a refined decomposition in Chow:

$$(1.9) \quad A^*(X) = \bigoplus_{i,s,\mu \in \mathfrak{t}^*} A^i(X)_{2s,\mu},$$

where (with $\lambda = (i - s)\tilde{h}^* + \mu$) we let

$$A^i(X)_{2s,\mu} = A^i(\mathfrak{h}_\lambda(X)) = \{\alpha \in A^i(X)_{2s} \mid h_v(\alpha) = \mu(v)\alpha \text{ for all } v \in \bar{\mathfrak{t}}\}.$$

Theorem 1.6. *As above, let $X = \text{Hilb}_n(S)$ with S a projective K3 surface.*

- (i) *The decomposition (1.8) is multiplicative, i.e., the cup product respects the weight decomposition, in the sense that*

$$\cup : \mathfrak{h}_\lambda(X) \otimes \mathfrak{h}_\mu(X) \rightarrow \mathfrak{h}_{\lambda+\mu}(X)$$

for all $\lambda, \mu \in \mathfrak{t}^$. As a result, the triple grading in (1.9) is multiplicative.*

- (ii) *All Chern classes of X belong to $A^*(X)_{0,0}$.*

By a result of Markman [22], $\bar{\mathfrak{g}}$ (namely the entire reduced LLV algebra, instead of just its Néron–Severi part; see [14]) is the Lie algebra of the monodromy group of X . Since the images of Chern classes in cohomology are invariant under monodromy, they are of weight 0 with respect to $\bar{\mathfrak{t}}$. We see that part (ii) of Theorem 1.6 confirms the expectation from the Beauville–Voisin conjecture.

The decompositions (1.8) and (1.9) can be defined also for abelian varieties starting from the decomposition (1.3) obtained by Moonen. The analogue of Theorem 1.6 (i) then follows from [24, Proposition 6.9 (iii)].

Since $\mathfrak{t} \subset \bar{\mathfrak{g}}_{\text{NS}} \oplus \mathbb{Q} \cdot \tilde{h}$, Theorem 1.6 (i) follows from the statement that any element of $\bar{\mathfrak{g}}_{\text{NS}}$ acts on $A^*(X)$ by derivations, akin to (1.6). The generators of $\bar{\mathfrak{g}}_{\text{NS}}$ are denoted by

$$h_{\alpha\beta} \in A^{2n}(X \times X)$$

and indexed by $\alpha \wedge \beta$ in $\wedge^2(A^1(X))$. We then deduce Theorem 1.6 (i) from the identity

$$(1.10) \quad h_{\alpha\beta}(x \cdot x') = h_{\alpha\beta}(x) \cdot x' + x \cdot h_{\alpha\beta}(x')$$

for all $\alpha, \beta \in A^1(X)$ (our proof of (1.10) will be at the level of correspondences; see Section 4).

It is natural to ask for an extension of our results to arbitrary hyper-Kähler varieties. By a result of Rieß [30], the Chow motives of two birational hyper-Kähler varieties are isomorphic as graded algebra objects. Moreover, the isomorphism preserves Chern classes. Hence our results here apply equally well to any hyper-Kähler variety birational to the Hilbert scheme of points of a K3 surface.

1.6. Conventions. Throughout the paper, Chow groups and Chow motives will be taken with \mathbb{Q} -coefficients. We refer to [25] for the definitions and conventions of Chow motives.

We will often switch between the languages of correspondences and operators on Chow groups, in the following sense. Every operator $f : A^*(X) \rightarrow A^*(Y)$ will arise from a correspondence $F \in A^*(X \times Y)$ by the usual construction

$$\begin{array}{ccc} & X \times Y & \\ \pi_1 \swarrow & & \searrow \pi_2 \\ X & & Y \end{array} \quad f = \pi_{2*}(F \cdot \pi_1^*)$$

and any compositions and equalities of operators implicitly entail compositions and equalities of correspondences. For example, the operator

$$\text{mult}_\tau : A^*(X) \rightarrow A^*(X)$$

of cup product with a fixed element $\tau \in A^*(X)$ is associated to the correspondence

$$\Delta_*(\tau) \in A^*(X \times X),$$

where $\Delta : X \hookrightarrow X \times X$ is the diagonal embedding.

Moreover, a family of operators $f_\gamma : A^*(X) \rightarrow A^*(Y)$ labeled by $\gamma \in A^*(Z)$ will arise from a correspondence $F \in A^*(X \times Y \times Z)$, by the assignment

$$f_\gamma \text{ arises from } \pi_{12*}(F \cdot \pi_3^*(\gamma)) \in A^*(X \times Y)$$

for all $\gamma \in A^*(Z)$. We employ the language of “operators indexed by $\gamma \in A^*(Z)$ ” instead of cycles on $X \times Y \times Z$ because it makes manifest the fact that γ does not play any role in taking compositions. For instance, the family of operators

$$\text{mult}_\gamma : A^*(X) \rightarrow A^*(X)$$

labeled by $\gamma \in A^*(X)$ is associated to the small diagonal $\Delta_{123} \subset X \times X \times X$.

We will often be concerned with cycles on a variety of the form $S^n = S \times \cdots \times S$ for a smooth algebraic variety S (most often an algebraic surface). We let

$$\Delta_{a_1 \dots a_k} \in A^*(S^n)$$

denote the diagonal $\{(x_1, \dots, x_n) \mid x_{a_1} = \cdots = x_{a_k}\}$, for all collections of distinct indices $a_1, \dots, a_k \in \{1, \dots, n\}$. Moreover, given a class $\Gamma \in A^*(S^k)$, we may choose to write it as $\Gamma_{1 \dots k}$ in order to indicate the power of S where this class lives. Then for any collection of distinct indices $a_1, \dots, a_k \in \{1, \dots, n\}$, we define

$$\Gamma_{a_1 \dots a_k} = p_{a_1 \dots a_k}^*(\Gamma) \in A^*(S^n),$$

where we let $p_{a_1 \dots a_k} = (p_{a_1}, \dots, p_{a_k}) : S^n \rightarrow S^k$ with $p_i : S^n \rightarrow S$ the projection to the i -th factor. Finally, if \bullet denotes any index from 1 to $k + 1$, we write

$$\int_{\bullet} : A^*(S^{k+1}) \rightarrow A^*(S^k)$$

for the push-forward map which forgets the factor labeled by \bullet .

2. Hilbert schemes

2.1. Throughout the paper, S will denote a projective K3 surface. In [3], Beauville and Voisin studied the class $c \in A^2(S)$ of any closed point on a rational curve in S , and they proved the following formulas in $A^*(S)$:

$$\begin{aligned} c_2(\text{Tan}_S) &= 24c, \\ \alpha \cdot \beta &= (\alpha, \beta)c \end{aligned}$$

for all $\alpha, \beta \in A^1(S)$ (above, we write $(\cdot, \cdot) : A^*(S) \otimes A^*(S) \rightarrow \mathbb{Q}$ for the intersection pairing). Moreover, we have the following identities in $A^*(S^2)$:

$$(2.1) \quad \Delta \cdot c_1 = \Delta \cdot c_2 = c_1 \cdot c_2,$$

$$(2.2) \quad \Delta \cdot \alpha_1 = \Delta \cdot \alpha_2 = \alpha_1 \cdot c_2 + \alpha_2 \cdot c_1,$$

where $\Delta \in A^*(S^2)$ is the class of the diagonal, and the following identity in $A^*(S^3)$:

$$(2.3) \quad \Delta_{123} = \Delta_{12} \cdot c_3 + \Delta_{13} \cdot c_2 + \Delta_{23} \cdot c_1 - c_1 \cdot c_2 - c_1 \cdot c_3 - c_2 \cdot c_3.$$

By iterating this identity, we obtain the corollary

$$(2.4) \quad \Delta_{1\dots k} = \sum_{1 \leq i < j \leq k} \Delta_{ij} \prod_{\ell \neq i, j} c_\ell - (k-2) \sum_{i=1}^k \prod_{\ell \neq i} c_\ell.$$

Proposition 2.1. *The following formulas hold:*

$$(2.5) \quad \gamma_1 c_1 = c_1 \int_{\bullet} \gamma_{\bullet} c_{\bullet}$$

$$(2.6) \quad \gamma_1 \alpha_1 = c_1 \int_{\bullet} \gamma_{\bullet} \alpha_{\bullet} + \alpha_1 \int_{\bullet} \gamma_{\bullet} c_{\bullet}$$

$$(2.7) \quad \gamma_1 \Delta_{1\dots k} = \sum_{i=1}^k \gamma_i \prod_{j \neq i} c_j + \left(\Delta_{1\dots k} - \sum_{i=1}^k \prod_{j \neq i} c_j \right) \int_{\bullet} c_{\bullet} \gamma_{\bullet} - (k-1) c_1 \dots c_k \int_{\bullet} \gamma_{\bullet}$$

for any $\gamma \in A^*(S \times S^l)$, where only the first index of γ appears in the equations above (the latter l indices are simply the same on the left and right-hand sides).

Proof. Formulas (2.5) and (2.6) both follow by taking (2.1) and (2.2) in $A^*(S \times S)$ (with the factors denoted by indices 1 and \bullet), multiplying them by γ_{\bullet} , and then integrating out the factor \bullet . As for (2.7), let us consider identity (2.3) in $A^*(S \times S \times S)$ (with the factors denoted by indices 1, 2, and \bullet) and multiply it by γ_{\bullet} . We obtain that

$$\Delta_{12\bullet} \gamma_{\bullet} = \Delta_{12c_{\bullet}} \gamma_{\bullet} + \Delta_{1\bullet c_2} \gamma_{\bullet} + \Delta_{2\bullet c_1} \gamma_{\bullet} - c_1 c_2 \gamma_{\bullet} - (c_1 + c_2) c_{\bullet} \gamma_{\bullet}$$

and hence

$$\Delta_{12\bullet} \gamma_1 = \Delta_{12c_{\bullet}} \gamma_{\bullet} + \Delta_{1\bullet c_2} \gamma_1 + \Delta_{2\bullet c_1} \gamma_2 - c_1 c_2 \gamma_{\bullet} - (c_1 + c_2) c_{\bullet} \gamma_{\bullet}.$$

If we integrate out the factor \bullet , we precisely obtain the $k = 2$ case of (2.7). To prove the general case of (2.7), we proceed by induction on k : the induction step is obtained by multiplying both sides of (2.7) with $\Delta_{k, k+1}$, and then applying (2.1), (2.2), and the $k = 2$ case of (2.7). \square

2.2. Consider the Hilbert scheme Hilb_n of n points on S and the Chow rings

$$\text{Hilb} = \bigsqcup_{n=0}^{\infty} \text{Hilb}_n, \quad A^*(\text{Hilb}) = \bigoplus_{n=0}^{\infty} A^*(\text{Hilb}_n)$$

always with rational coefficients. We will consider two types of elements of the Chow rings above. The first of these are defined by considering the universal subscheme

$$\mathcal{Z}_n \subset \text{Hilb}_n \times S.$$

For any $k \in \mathbb{N}$, consider the projections

$$\text{Hilb}_n \xleftarrow{\pi} \text{Hilb}_n \times S^k \xrightarrow{\rho} S^k$$

and let $\mathcal{Z}_n^{(i)} \subset \text{Hilb}_n \times S^k$ denote the pullback of \mathcal{Z}_n via the i -th projection $S^k \rightarrow S$.

Definition 2.2. A *universal class* is any element of $A^*(\text{Hilb}_n)$ of the form

$$(2.8) \quad \pi_* c [P(\dots, \text{ch}_j(\mathcal{O}_{\mathcal{Z}_n^{(i)}}), \dots)_{j \in \mathbb{N}}^{1 \leq i \leq k} \pi]$$

for all $k \in \mathbb{N}$ and for all polynomials P with coefficients pulled back from $A^*(S^k)$.

In particular, by inserting diagonals if necessary, the classes (2.8) where P is a monomial are of the form

$$(2.9) \quad \text{univ}_{d_1, \dots, d_k}(\Gamma) = \pi_* \pi [\text{ch}_{d_1}(\mathcal{O}_{\mathcal{Z}_n^{(1)}}) \dots \text{ch}_{d_k}(\mathcal{O}_{\mathcal{Z}_n^{(k)}}) \cdot \rho^*(\Gamma) \pi].$$

The following theorem holds for every smooth quasi-projective surface (see [28]), but we only prove it here in the case where S is a K3 surface (the argument herein easily generalizes to any smooth projective surface using the results of [13]).

Theorem 2.3. Any class in $A^*(\text{Hilb}_n)$ is universal, i.e., of the form (2.8).

Proof. Consider the product $\text{Hilb}_n \times S^k \times \text{Hilb}_n$, and we will write $\pi_1, \pi_2, \pi_3, \pi_{12}, \pi_{23}$, and π_{13} for the various projections to its factors. As a consequence of [21] (see also [13]), the diagonal $\Delta_{\text{Hilb}_n} \subset \text{Hilb}_n \times \text{Hilb}_n$ can be written as follows:

$$\Delta_{\text{Hilb}_n} = \pi_{13*} \left[\sum_a \pi_2^*(\gamma_a) \prod_{(i,j)} \text{ch}_j(\mathcal{O}_{\mathcal{Z}_n^{(i)}}) \prod_{(\tilde{i}, \tilde{j})} \text{ch}_{\tilde{j}}(\mathcal{O}_{\tilde{\mathcal{Z}}_n^{(\tilde{i})}}) \right]$$

for suitably chosen $k \in \mathbb{N}$, where we do not care much about the specific coefficients γ_a and indices $i, j, \tilde{i}, \tilde{j}$ which appear in the sum above (we write \mathcal{Z}_n and $\tilde{\mathcal{Z}}_n$ for the universal subschemes in $\text{Hilb}_n \times S \times \text{Hilb}_n$ corresponding to the first and second copies of Hilb_n , respectively). Since the diagonal corresponds to the identity operator, the equality above implies that

$$(2.10) \quad \begin{aligned} \text{Id}_{\text{Hilb}_n} &= \pi_{1*} \left[\sum_a \pi_2^*(\gamma_a) \prod_{(i,j)} \text{ch}_j(\mathcal{O}_{\mathcal{Z}_n^{(i)}}) \prod_{(\tilde{i}, \tilde{j})} \text{ch}_{\tilde{j}}(\mathcal{O}_{\tilde{\mathcal{Z}}_n^{(\tilde{i})}}) \pi_3^* \right] \\ &= \sum_a \pi_* \left[\prod_{(i,j)} \text{ch}_j(\mathcal{O}_{\mathcal{Z}_n^{(i)}}) \rho^* \left(\gamma_a \cdot \rho_* \left(\prod_{(\tilde{i}, \tilde{j})} \text{ch}_{\tilde{j}}(\mathcal{O}_{\tilde{\mathcal{Z}}_n^{(\tilde{i})}}) \cdot \pi^* \right) \right) \right] \end{aligned}$$

hence the universality. \square

Formula (2.10) implies the surjectivity of the homomorphism

$$(2.11) \quad \bigoplus_a A^*(S^k) \twoheadrightarrow A^*(\text{Hilb}_n), \quad \sum_a \Gamma_a \mapsto \sum_a \pi_* \left[\prod_{(i,j)} \text{ch}_j(\mathcal{O}_{Z_n^{(i)}}) \rho^*(\Gamma_a) \right],$$

where the sums over a are in one-to-one correspondence with the sums in (2.10).

2.3. Let us present another important source of elements of $A^*(\text{Hilb}_n)$, based on the following construction independently due to Grojnowski [15] and Nakajima [26] (in the present paper, we will mostly use the presentation by Nakajima). For any $n, k \in \mathbb{N}$, consider the closed subscheme

$$\text{Hilb}_{n,n+k} = \{(I \supset I') \mid I/I' \text{ is supported at a single } x \in S\} \subset \text{Hilb}_n \times \text{Hilb}_{n+k}$$

endowed with projection maps

$$\begin{array}{ccc} & \text{Hilb}_{n,n+k} & \\ p_- \swarrow & \downarrow p_S & \searrow p_+ \\ \text{Hilb}_n & S & \text{Hilb}_{n+k} \end{array}$$

that remember I, x, I' , respectively. One may use $\text{Hilb}_{n,n+k}$ as a correspondence

$$(2.12) \quad A^*(\text{Hilb}_n) \xrightarrow{\mathfrak{q}_{\pm k}} A^*(\text{Hilb}_{n \pm k} \times S)$$

given by

$$(2.13) \quad \mathfrak{q}_{\pm k} = (\pm 1)^k \cdot (p_{\pm} \times p_S)_* \circ p_{\mp}^*.$$

Because the correspondences above are defined for all n , it makes sense to set

$$A^*(\text{Hilb}) \xrightarrow{\mathfrak{q}_{\pm k}} A^*(\text{Hilb} \times S).$$

We also set $\mathfrak{q}_0 = 0$. The main result of [26] (although [26] is written at the level of cohomology, the result holds at the level of Chow groups; see for example [27, Remark 8.15 (2)]) is that the operators \mathfrak{q}_k obey the commutation relations in the Heisenberg algebra, namely

$$(2.14) \quad [\mathfrak{q}_k, \mathfrak{q}_l] = k \delta_{k+l}^0 (\text{Id}_{\text{Hilb}} \times \Delta)$$

as correspondences

$$A^*(\text{Hilb}) \rightarrow A^*(\text{Hilb} \times S^2).$$

In terms of self-correspondences $A^*(\text{Hilb}) \rightarrow A^*(\text{Hilb})$, the identity in equation (2.14) reads, for all $\alpha, \beta \in A^*(S)$,

$$(2.15) \quad [\mathfrak{q}_k(\alpha), \mathfrak{q}_l(\beta)] = k(\alpha, \beta) \text{Id}_{\text{Hilb}}.$$

2.4. More generally, we may consider

$$(2.16) \quad \mathfrak{q}_{n_1} \dots \mathfrak{q}_{n_t} : A^*(\text{Hilb}) \rightarrow A^*(\text{Hilb} \times S^t),$$

where the convention is that the operator \mathfrak{q}_{n_i} acts in the i -th factor of $S^t = S \times \dots \times S$. Then

associated to any $\Gamma \in A^*(S^t)$, one obtains an endomorphism of $A^*(\text{Hilb})$:

$$(2.17) \quad \mathfrak{q}_{n_1} \cdots \mathfrak{q}_{n_t}(\Gamma) = \pi_*(\rho^*(\Gamma) \cdot \mathfrak{q}_{n_1} \cdots \mathfrak{q}_{n_t})$$

where π and ρ denote the projections of $\text{Hilb} \times S^t$ to the factors.

Theorem 2.4 ([4]). *We have a decomposition*

$$(2.18) \quad A^*(\text{Hilb}) = \bigoplus_{\substack{n_1 \geq \cdots \geq n_t \in \mathbb{N} \\ \Gamma \in A^*(S^t)^{\text{sym}}}} \mathfrak{q}_{n_1} \cdots \mathfrak{q}_{n_t}(\Gamma) \cdot v,$$

where “sym” refers to the part of $A^*(S^t)$ which is symmetric with respect to those transpositions $(ij) \in \mathfrak{S}_t$ for which $n_i = n_j$, and v is a generator of $A^*(\text{Hilb}_0) \cong \mathbb{Q}$.

Proof. Since we will need it later, we recall the precise relationship between Nakajima operators and the correspondences studied in [4]. Let λ be a partition of n with k parts, let $S^\lambda = S^k$ and let $S^\lambda \rightarrow S^{(n)}$ be the map that sends (x_1, \dots, x_k) to the cycle $\lambda_1 x_1 + \cdots + \lambda_k x_k$ in the n -th symmetric product of the surface S . We consider the correspondence

$$\Gamma_\lambda = (\text{Hilb}_n \times_{S^{(n)}} S^\lambda)_{\text{red}} = \{(I, x_1, \dots, x_k) \mid \sigma(I) = \lambda_1 x_1 + \cdots + \lambda_k x_k\},$$

where $\sigma : S^{[n]} \rightarrow S^{(n)}$ is the Hilbert–Chow morphism sending the subscheme I to its underlying support. The subscheme Γ_λ is irreducible of dimension $n + k$ and the locus $\Gamma_\lambda^{\text{reg}} \subset \Gamma_\lambda$, where the points x_i are distinct, is open and dense; see [4, Remark 2.0.1]. Similarly, the Nakajima correspondence $\mathfrak{q}_{\lambda_1} \cdots \mathfrak{q}_{\lambda_k}$ is a cycle in $\text{Hilb}_n \times S^k$ of dimension $n + k$ supported on a subscheme that contains $\Gamma_\lambda^{\text{reg}}$ as an open subset and whose complement is of smaller dimension [26, Section 4(i)]. Moreover, the multiplicity of the cycle on $\Gamma_\lambda^{\text{reg}}$ is 1. Hence we have the equality of correspondences

$$(2.19) \quad \Gamma_\lambda = \mathfrak{q}_{\lambda_1} \cdots \mathfrak{q}_{\lambda_k} \in A^*(\text{Hilb}_n \times S^\lambda).$$

The result follows now from [4, Proposition 6.1.5], which says that

$$(2.20) \quad \Delta_{\text{Hilb}_n} = \sum_{\lambda \vdash n} \frac{(-1)^{n-l(\lambda)}}{|\text{Aut}(\lambda)| \prod_i \lambda_i} {}^t \Gamma_\lambda \circ \Gamma_\lambda$$

where λ runs over all partitions of size n , and we let $l(\lambda)$ and λ_i denote the length and the parts of λ , respectively. \square

Remark 2.5. As shown in [28], there is an explicit way to go between the descriptions (2.8) and (2.18) of $A^*(\text{Hilb})$. Concretely, for all $n_1 \geq \cdots \geq n_t$ there exists a polynomial P_{n_1, \dots, n_t} with coefficients in $\rho^*(A^*(S^t))$ such that for all $\Gamma \in A^*(S^t)$,

$$\mathfrak{q}_{n_1} \cdots \mathfrak{q}_{n_t}(\Gamma) = \pi_* [P_{n_1, \dots, n_t}(\dots, \text{ch}_j(\mathcal{O}_{\mathbb{Z}^{(i)}}), \dots)_{j \in \mathbb{N}}^{1 \leq i \leq t} \cdot \rho^*(\Gamma)].$$

Moreover, [28] gives an algorithm for computing the polynomial P_{n_1, \dots, n_t} .

2.5. Two interesting collections of elements of $A^*(\text{Hilb}_n)$ can be written as universal classes: divisors and Chern classes of the tangent bundle. One has

$$A^1(S) \oplus \mathbb{Q} \cdot \delta \cong A^1(\text{Hilb}_n)$$

(with the convention that $\delta = 0$ if $n = 1$), where the isomorphism is given by

$$\begin{aligned} l \in A^1(S) &\mapsto \text{univ}_2(l), \\ \delta &\mapsto \text{univ}_3(1). \end{aligned}$$

Similarly, the Chern character of the tangent bundle to Hilb_n is given by the well-known formula (see for example [23, Proposition 2.10])

$$\text{ch}(\text{Tan}_{\text{Hilb}_n}) = \pi_* \left[(\text{ch}(\mathcal{O}_{\mathbb{Z}^n}) + \text{ch}(\mathcal{O}_{\mathbb{Z}^n})' - \text{ch}(\mathcal{O}_{\mathbb{Z}^n})\text{ch}(\mathcal{O}_{\mathbb{Z}^n})') \rho^*(1 + 2c) \right],$$

where $(\cdot)'$ is the operator which multiplies a degree d class by $(-1)^d$. Therefore, the Chern character of the tangent bundle is a linear combination of the following particular universal classes:

$$\text{univ}_d(\gamma) \quad \text{and} \quad \text{univ}_{d,d'}(\Delta_*(\gamma)),$$

where $\gamma \in \{1, c\}$, and d, d' are various natural numbers.

3. Motivic decompositions

3.1. Recall the Lie algebra action $\mathfrak{g}_{\text{NS}} \curvearrowright A^*(\text{Hilb}_n)$ from [29], which lifts the classical construction of [20, 34] in cohomology. To this end, consider the Beauville–Bogomolov form, which is the pairing on

$$V = A^1(\text{Hilb}_n) \cong A^1(S) \oplus \mathbb{Q} \cdot \delta.$$

This form extends the intersection form on $A^1(S)$ and satisfies

$$(\delta, \delta) = 2 - 2n, \quad (\delta, A^1(S)) = 0.$$

Let $U = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ be the hyperbolic lattice with fixed symplectic basis e, f . We have

$$\mathfrak{g}_{\text{NS}} = \wedge^2(V \oplus U_{\mathbb{Q}}),$$

where the Lie bracket is defined for all $a, b, c, d \in V \oplus U_{\mathbb{Q}}$ by

$$[a \wedge b, c \wedge d] = (a, d)b \wedge c - (a, c)b \wedge d - (b, d)a \wedge c + (b, c)a \wedge d.$$

Consider for all $\alpha \in A^1(S)$ the following operators:

$$\begin{aligned} (3.1) \quad e_{\alpha} &= - \sum_{n>0} \mathfrak{q}_n \mathfrak{q}_{-n}(\Delta_* \alpha), \\ e_{\delta} &= -\frac{1}{6} \sum_{i+j+k=0} : \mathfrak{q}_i \mathfrak{q}_j \mathfrak{q}_k (\Delta_{123}) :, \\ \tilde{f}_{\alpha} &= - \sum_{n>0} \frac{1}{n^2} \mathfrak{q}_n \mathfrak{q}_{-n}(\alpha_1 + \alpha_2), \\ \tilde{f}_{\delta} &= -\frac{1}{6} \sum_{i+j+k=0} : \mathfrak{q}_i \mathfrak{q}_j \mathfrak{q}_k \left(\frac{1}{k^2} \Delta_{12} + \frac{1}{j^2} \Delta_{13} + \frac{1}{i^2} \Delta_{23} \right. \\ &\quad \left. + \frac{2}{jk} c_1 + \frac{2}{ik} c_2 + \frac{2}{ij} c_3 \right) :. \end{aligned}$$

Here $:-:$ is the normal ordered product defined by

$$(3.2) \quad :q_{i_1} \dots q_{i_k} := q_{i_{\sigma(1)}} \dots q_{i_{\sigma(k)}},$$

where σ is any permutation such that $i_{\sigma(1)} \geq \dots \geq i_{\sigma(k)}$. We define operators e_α and \widetilde{f}_α for general $\alpha \in A^1(\text{Hilb}_n)$ by linearity in α . By [23, Theorem 1.6], we have that e_α is the operator of cup product with α . If $(\alpha, \alpha) \neq 0$, the multiple $\widetilde{f}_\alpha/(\alpha, \alpha)$ acts on cohomology as the Lefschetz dual of e_α . In [29], it was shown that the assignment

$$(3.3) \quad \text{act} : \mathfrak{g}_{\text{NS}} \rightarrow A^*(\text{Hilb}_n \times \text{Hilb}_n), \quad \text{act}(e \wedge \alpha) = e_\alpha, \quad \text{act}(\alpha \wedge f) = \widetilde{f}_\alpha$$

for all $\alpha \in V$, induces a Lie algebra homomorphism. In particular, it was shown in [29] that the element $e \wedge f \in \mathfrak{g}_{\text{NS}}$ acts by

$$(3.4) \quad h = \sum_{k>0} \frac{1}{k} q_k q_{-k} (c_2 - c_1).$$

The operator h specializes in cohomology to the Lefschetz grading operator, which by our normalization acts on the space $H^{2i}(\text{Hilb}_n)$ by multiplication by $i - n$. Similarly, for any $\alpha, \beta \in A^1(S) \subset A^1(\text{Hilb}_n)$, the element $\alpha \wedge \beta \in \mathfrak{g}_{\text{NS}}$ acts by

$$(3.5) \quad h_{\alpha\beta} = \sum_{k=1}^{\infty} \frac{1}{k} q_k q_{-k} (\alpha_2 \beta_1 - \alpha_1 \beta_2)$$

and the element $\alpha \wedge \delta \in \mathfrak{g}_{\text{NS}}$ acts by

$$(3.6) \quad h_{\alpha\delta} = -\frac{1}{2} \sum_{\substack{i+j+k=0 \\ i,j,k \in \mathbb{Z}}} \frac{1}{k} :q_i q_j q_k (\Delta_{12}(\alpha_1 + \alpha_3)):$$

3.2. Proof of Theorem 1.3. Let us start with the decomposition of the diagonal into Nakajima operators

$$(3.7) \quad \Delta_{\text{Hilb}_n} = \sum_{\lambda \vdash n} \frac{(-1)^{l(\lambda)}}{\mathfrak{z}(\lambda)} q_\lambda q_{-\lambda}(\Delta),$$

where λ runs over all partitions of n ,

$$\mathfrak{z}(\lambda) = |\text{Aut}(\lambda)| \prod_i \lambda_i$$

is a combinatorial factor, and for any $\pi \in A^*(S \times S)$ we write

$$\begin{aligned} q_\lambda q_{-\lambda}(\pi) &= q_{\lambda_1} \dots q_{\lambda_{l(\lambda)}} q_{-\lambda_1} \dots q_{-\lambda_{l(\lambda)}} (\pi_{1,l(\lambda)+1} \pi_{2,l(\lambda)+2} \dots \pi_{l(\lambda),2l(\lambda)}) \\ &= :q_{\lambda_1} q_{-\lambda_1}(\pi) \dots q_{\lambda_{l(\lambda)}} q_{-\lambda_{l(\lambda)}}(\pi):. \end{aligned}$$

Formula (3.7) follows directly from (2.19), (2.20), and the fact that ${}^t q_m = (-1)^m q_{-m}$ (which is incorporated in the definition (2.13)).

Consider the decomposition of the diagonal of S as

$$(3.8) \quad \Delta = \pi_{-1} + \pi_0 + \pi_1,$$

where

$$\pi_{-1} = c_1, \quad \pi_0 = \Delta - c_1 - c_2, \quad \pi_1 = c_2.$$

It is easy to note that π_{-1}, π_0, π_1 are the projectors onto the $-1, 0, +1$ -eigenspaces of the action of h on $A^*(\text{Hilb}_1) = A^*(S)$.

To define projectors corresponding to the action of h on $A^*(\text{Hilb}_n)$, we insert the decomposition (3.8) into (3.7), and then expand and collect the terms of degree i . Concretely, for every integer i , we let

$$P_i = \sum_{\substack{\lambda, \mu, \nu \\ |\lambda|+|\mu|+|\nu|=n \\ -l(\lambda)+l(\nu)=i}}^{l(\lambda), l(\mu), l(\nu)} \frac{(-1)^{l(\lambda)+l(\mu)+l(\nu)}}{\mathfrak{z}(\lambda)\mathfrak{z}(\mu)\mathfrak{z}(\nu)} : \mathfrak{q}_\lambda \mathfrak{q}_{-\lambda}({}^t \pi_{-1}) \mathfrak{q}_\mu \mathfrak{q}_{-\mu}({}^t \pi_0) \mathfrak{q}_\nu \mathfrak{q}_{-\nu}({}^t \pi_1) :.$$

In particular, we have $P_i = 0$ unless $i \in \{-n, \dots, n\}$. Let us check that P_i are indeed projectors onto the eigenspaces of h .

Claim 3.1. *For all integers $i, j \in \{-n, \dots, n\}$, we have the following equalities in the ring $A^*(\text{Hilb}_n \times \text{Hilb}_n)$:*

- (a) $P_i \circ P_j = P_i \delta_j^i$,
- (b) $h \circ P_i = i P_i$.

Proof. (a) We determine $P_i \circ P_j$ by commuting all Nakajima operators with negative indices to the right, and then using that we act on Hilb_n so all products of Nakajima operators with purely negative indices of degree $> n$ vanish. Since every summand in P_j contains such a product of degree n , we find that for a term to contribute all operators with negative indices coming from P_i have to interact with operators (with positive indices) from the second term. The interactions are described as follows. For a single term (let $a, b > 0$ and $r, s \in \{-1, 0, 1\}$) we have

$$\begin{aligned} \mathfrak{q}_a \mathfrak{q}_{-a}({}^t \pi_r) \mathfrak{q}_b \mathfrak{q}_{-b}({}^t \pi_s) &= \mathfrak{q}_a [\mathfrak{q}_{-a}, \mathfrak{q}_b] \mathfrak{q}_{-b}({}^t \pi_r) {}_{12}({}^t \pi_s)_{34} \\ &\quad + \mathfrak{q}_a \mathfrak{q}_b \mathfrak{q}_{-a} \mathfrak{q}_{-b}({}^t \pi_r) {}_{13}({}^t \pi_s)_{24}, \end{aligned}$$

where by the commutation relations (2.14) the first term on the right is

$$\begin{aligned} \mathfrak{q}_a [\mathfrak{q}_{-a}, \mathfrak{q}_b] \mathfrak{q}_{-b}({}^t \pi_r) {}_{12}({}^t \pi_s)_{34} &= (-a) \delta_{ab} \mathfrak{q}_a \mathfrak{q}_{-b} (\pi_{14} * ({}^t \pi_r) {}_{12}({}^t \pi_s)_{34} \Delta_{23}) \\ &= (-a) \delta_{ab} \mathfrak{q}_a \mathfrak{q}_{-a} ({}^t \pi_s \circ {}^t \pi_r) \\ &= (-a) \delta_{ab} \mathfrak{q}_a \mathfrak{q}_{-a} ({}^t (\pi_r \circ \pi_s)) \\ &= (-a) \delta_{ab} \delta_{rs} \mathfrak{q}_a \mathfrak{q}_{-a} ({}^t \pi_r). \end{aligned}$$

Hence for a composition

$$: \mathfrak{q}_\lambda \mathfrak{q}_{-\lambda}({}^t \pi_{-1}) \mathfrak{q}_\mu \mathfrak{q}_{-\mu}({}^t \pi_0) \mathfrak{q}_\nu \mathfrak{q}_{-\nu}({}^t \pi_1) : \circ : \mathfrak{q}_{\lambda'} \mathfrak{q}_{-\lambda'}({}^t \pi_{-1}) \mathfrak{q}_{\mu'} \mathfrak{q}_{-\mu'}({}^t \pi_0) \mathfrak{q}_{\nu'} \mathfrak{q}_{-\nu'}({}^t \pi_1) :$$

(with λ, μ, ν as in the definition of P_i , and the same for the primed partitions) to act non-trivially on Hilb_n we have to have $\lambda = \lambda'$, $\mu = \mu'$ and $\nu = \nu'$. Moreover, if we write λ multiplicatively as $(1^{l_1} 2^{l_2} \dots)$, where l_i is the number of parts of size i , there are precisely $|\text{Aut}(\lambda)| = \prod_i l_i!$ different ways to pair the negative factors in $\mathfrak{q}_\lambda \mathfrak{q}_{-\lambda}({}^t \pi_{-1})$ with the positive factors $\mathfrak{q}_{\lambda'} \mathfrak{q}_{-\lambda'}({}^t \pi_{-1})$, and similarly for μ, ν . Hence

$$\begin{aligned} : \mathfrak{q}_\lambda \mathfrak{q}_{-\lambda}({}^t \pi_{-1}) \mathfrak{q}_\mu \mathfrak{q}_{-\mu}({}^t \pi_0) \mathfrak{q}_\nu \mathfrak{q}_{-\nu}({}^t \pi_1) : \circ : \mathfrak{q}_{\lambda'} \mathfrak{q}_{-\lambda'}({}^t \pi_{-1}) \mathfrak{q}_{\mu'} \mathfrak{q}_{-\mu'}({}^t \pi_0) \mathfrak{q}_{\nu'} \mathfrak{q}_{-\nu'}({}^t \pi_1) : \\ = \delta_{\lambda \lambda'} \delta_{\mu \mu'} \delta_{\nu \nu'} (-1)^{l(\lambda)+l(\mu)+l(\nu)} \mathfrak{z}(\lambda) \mathfrak{z}(\mu) \mathfrak{z}(\nu) : \mathfrak{q}_\lambda \mathfrak{q}_{-\lambda}({}^t \pi_{-1}) \mathfrak{q}_\mu \mathfrak{q}_{-\mu}({}^t \pi_0) \mathfrak{q}_\nu \mathfrak{q}_{-\nu}({}^t \pi_1) : \end{aligned}$$

which implies the claim.

(b) To determine $h \circ P_i$ we commute h into the middle, i.e., to the right of all Nakajima operators with positive indices, and to the left of all with negative ones. In the middle position h acts on the Chow ring of Hilb_0 , where it vanishes. Hence again we only need to compute the commutators. For this we use (4.19) and that π_i are the projectors onto the eigenspaces of h so that

$$(h \times \text{Id})({}^t \pi_r) = {}^t((\text{Id} \times h)(\pi_r)) = {}^t(h \circ \pi_r) = r({}^t \pi_r).$$

As desired we find

$$h \circ P_i = (-1 \cdot l(\lambda) + 0 \cdot l(\mu) + 1 \cdot l(\nu))P_i = iP_i. \quad \square$$

By using the claim, it follows that the motivic decomposition:

$$\mathfrak{h}(\text{Hilb}_n) = \bigoplus_{i=0}^{2n} \mathfrak{h}^{2i}(\text{Hilb}_n)$$

with $\mathfrak{h}^{2i}(\text{Hilb}_n) = (\text{Hilb}_n, P_{i-n})$ has the stated properties. The uniqueness of the decomposition follows from the uniqueness of the decomposition of Δ_{Hilb_n} under the action of h on $A^*(\text{Hilb}_n \times \text{Hilb}_n)$; see the proof for the refined decomposition in Section 3.3 below. \square

By (2.20), an alternative way to write the projector P_i is

$$P_i = \sum_{\lambda \vdash n} \frac{(-1)^{n-l(\lambda)}}{\mathfrak{z}(\lambda)} {}^t \Gamma_\lambda \circ \tilde{P}_i \circ \Gamma_\lambda,$$

where $\tilde{P}_i \in A^*(S^\lambda \times S^\lambda)$ is the projector

$$\tilde{P}_i = \sum_{i_1 + \dots + i_{l(\lambda)} = i} \pi_{i_1} \times \dots \times \pi_{i_{l(\lambda)}}.$$

Hence the decomposition of Theorem 1.3 is precisely the Chow–Künneth decomposition constructed by Vial in [35, Section 2].

3.3. Refined decomposition. Let $U(\mathfrak{g}_{\text{NS}})$ be the universal enveloping algebra of \mathfrak{g}_{NS} . The Lie algebra homomorphism (3.3) extends to an algebra homomorphism

$$\text{act} : U(\mathfrak{g}_{\text{NS}}) \rightarrow A^*(\text{Hilb}_n \times \text{Hilb}_n).$$

Lemma 3.2. *The image $W \subset A^*(\text{Hilb}_n \times \text{Hilb}_n)$ of act is finite-dimensional.*

Proof. For every fixed $k \geq 1$ the subring of $R^*(S^k) \subset A^*(S^k)$ generated by

- α_i for all i and $\alpha \in A^1(S)$,
- c_i for all i ,
- Δ_{ij} for all i, j

is finite-dimensional, and preserved by the projections to the factors. Hence the space of operators $\tilde{W} \subset A^*(\text{Hilb}_n \times \text{Hilb}_n)$ spanned by

$$\mathfrak{q}_{\lambda_1} \cdots \mathfrak{q}_{\lambda_{l(\lambda)}} \mathfrak{q}_{-\mu_1} \cdots \mathfrak{q}_{-\mu_{l(\mu)}}(\Gamma)$$

for all partitions λ, μ of n and all $\Gamma \in R^*(S^{l(\lambda)+l(\mu)})$ is finite-dimensional. The commutation relations (2.14) show that \widetilde{W} is closed under compositions of correspondences. Moreover, by inspecting the expressions for the generators of \mathfrak{g}_{NS} in (3.1) (and using (3.7) to bring them into the desired form), we see that all generators of \mathfrak{g}_{NS} lie in \widetilde{W} . Hence $g \in \widetilde{W}$ for all $g \in U(\mathfrak{g}_{\text{NS}})$, i.e., $W \subset \widetilde{W}$. \square

We find that W is a finite-dimensional vector space which is preserved by the action of $U(\mathfrak{g}_{\text{NS}})$, and hence defines a finite-dimensional representation of \mathfrak{g}_{NS} . Since \mathfrak{g}_{NS} is semi-simple, this representation decomposes into isotypic summands

$$W = \bigoplus_{\psi \in \text{Irrep}(\mathfrak{g}_{\text{NS}})} W_{\psi}.$$

Let us look at the image of $\Delta_{\text{Hilb}_n} \in W$ under this decomposition:

$$(3.9) \quad \Delta_{\text{Hilb}_n} = \sum_{\psi \in \text{Irrep}(\mathfrak{g}_{\text{NS}})} P_{\psi},$$

where $P_{\psi} \in W_{\psi}$.

Claim 3.3. *The elements $P_{\psi} \in A^*(\text{Hilb}_n \times \text{Hilb}_n)$ are orthogonal projectors.*

Proof. Let us first show that left-multiplication by P_{ψ} maps W to W_{ψ} , i.e.,

$$(3.10) \quad P_{\psi} \circ W \subset W_{\psi}.$$

Indeed, for all $a \in W$, right multiplication by a is a \mathfrak{g}_{NS} -intertwiner and thus sends W_{ψ} to W_{ψ} . In other words, we have $W_{\psi} \circ a \subset W_{\psi}$, hence $W_{\psi} \circ W \subset W_{\psi}$, which implies (3.10). If we multiply any $a \in W$ by relation (3.9), we obtain

$$a = \sum_{\psi \in \text{Irrep}(\mathfrak{g}_{\text{NS}})} P_{\psi} \circ a.$$

By (3.10), the summands in the right-hand side each lie in W_{ψ} . If $a \in W_{\psi'}$, then by comparing summands the equality above implies

$$P_{\psi'} \circ a = a \quad \text{and} \quad P_{\psi} \circ a = 0$$

for all $\psi \neq \psi'$. In particular, taking $a = P_{\psi'}$ implies the relations $P_{\psi} \circ P_{\psi'} = \delta_{\psi\psi'} P_{\psi}$. Moreover, this implies that the inclusion (3.10) is actually an identity, hence left multiplication by P_{ψ} projects W onto W_{ψ} . \square

From Claim 3.3 we obtain the decomposition

$$(3.11) \quad \mathfrak{h}(\text{Hilb}_n) = \bigoplus_{\psi \in \text{Irrep}(\mathfrak{g}_{\text{NS}})} \mathfrak{h}_{\psi}(\text{Hilb}_n),$$

where $\mathfrak{h}_{\psi}(\text{Hilb}_n) = (\text{Hilb}_n, P_{\psi})$. We can now prove the main result of this section.

Proof of Theorem 1.5. It remains to show that the summands $\mathfrak{h}_{\psi}(\text{Hilb}_n)$ are ψ -isotypic and that the decomposition (3.11) is unique. Let M be a Chow motive. The action of \mathfrak{g}_{NS} on $\text{Hom}(M, \mathfrak{h}(\text{Hilb}_n))$ is defined by $g \mapsto \text{act}(g) \circ (-)$. Hence if $f \in \text{Hom}(M, M')$ is a morphism of Chow motives, the pullback

$$f^* : \text{Hom}(M', \mathfrak{h}(\text{Hilb}_n)) \rightarrow \text{Hom}(M, \mathfrak{h}(\text{Hilb}_n))$$

is equivariant with respect to the \mathfrak{g}_{NS} -action. Now, for any $v \in \text{Hom}(M, \mathfrak{h}_\psi(\text{Hilb}_n))$ we have $v = P_\psi \circ w$ for some $w \in \text{Hom}(M, \mathfrak{h}(\text{Hilb}_n))$ and thus

$$U(\mathfrak{g}_{\text{NS}})v = U(\mathfrak{g}_{\text{NS}})w^*(P_\psi) = w^*(U(\mathfrak{g}_{\text{NS}}) \circ P_\psi).$$

Since $U(\mathfrak{g}_{\text{NS}}) \circ P_\psi \subset W_\psi$, this implies that $U(\mathfrak{g}_{\text{NS}})v$ is finite-dimensional and ψ -isotypic. Since v was arbitrary, we conclude that $\text{Hom}(M, \mathfrak{h}_\psi(\text{Hilb}_n))$ is ψ -isotypic.

The decomposition (3.11) is unique because (3.9) is unique. Indeed, suppose we had any other decomposition

$$(3.12) \quad \Delta_{\text{Hilb}_n} = \sum_{\psi \in \text{Irrrep}(\mathfrak{g}_{\text{NS}})} P'_\psi,$$

where $P'_\psi \in W_\psi$ for all ψ . Then we would need $P'_\psi = P_\psi \circ a_\psi$ for some $a_\psi \in W$. But multiplying (3.12) on the left with P_ψ and using the orthogonality of the projectors would imply $P_\psi = P_\psi \circ P_\psi \circ a_\psi = P_\psi \circ a_\psi = P'_\psi$. \square

As in [24, Proof of Theorem 7.2], we could also have used Yoneda's Lemma to conclude the existence of the decomposition (3.11). Our presentation above has the advantage of being constructive. It also shows that the projectors P_ψ can be written in terms of the Nakajima operators applied to elements in $R^*(S^k)$.

4. Multiplicativity

4.1. The main purpose of the present section is to prove Theorems 1.4 and 1.6. Let us first discuss the general strategy. Given an operator $H : A^*(\text{Hilb}_n) \rightarrow A^*(\text{Hilb}_n)$ among $h, h_{\alpha\beta}, h_{\alpha\delta}$, we will first prove a commutation relation of the form

$$(4.1) \quad [H, \text{mult}_x] = \text{mult}_y,$$

where mult_x is the operator of multiplication by any $x \in A^*(\text{Hilb}_n)$ and y will be given by an explicit formula in terms of x . This equation will help us in two ways: Firstly, applying (4.1) to the fundamental class $1_n \in A^0(\text{Hilb}_n)$ yields

$$(4.2) \quad H(x) - H(1_n)x = y.$$

Hence if we define $\tilde{H} = H - H(1_n)\text{Id}_{\text{Hilb}_n}$, then (4.1) reads

$$[\tilde{H}, \text{mult}_x] = \text{mult}_{\tilde{H}(x)}.$$

In other words, we have $\tilde{H}(x \cdot x') = \tilde{H}(x) \cdot x' + x \cdot \tilde{H}(x')$ for all x, x' , which precisely states that \tilde{H} is multiplicative. Secondly, the explicit formula for y together with (4.2) will yield an expression for $H(x)$, namely $y + H(1_n)x$. This will be used to determine the value of H on Chern and divisor classes.

4.2. In order to prove relations of the form (4.1), we will introduce the machinery of operators on Chow groups developed by [18, 19, 23]. The main idea is to develop a common framework for studying the Nakajima operators (2.12) and the following operators:

$$(4.3) \quad \mathcal{G}_d : A^*(\text{Hilb}) \xrightarrow{\pi^*} A^*(\text{Hilb} \times S) \xrightarrow{\text{mult}_{\text{ch}_d(\mathcal{O}_{\mathbb{Z}})}} A^*(\text{Hilb} \times S)$$

(where $\pi : \text{Hilb} \times S^t \rightarrow \text{Hilb}$ denotes the first projection). We will employ the following notation for compositions of these operators, akin to (2.16) and (2.17):

$$\mathfrak{G}_{d_1} \dots \mathfrak{G}_{d_t} : A^*(\text{Hilb}) \rightarrow A^*(\text{Hilb} \times S^t),$$

where the convention is that \mathfrak{G}_{d_i} acts on the i -th factor of $S^t = S \times \dots \times S$. Then associated to any $\Gamma \in A^*(S^t)$, one obtains the following endomorphism:

$$\mathfrak{G}_{d_1} \dots \mathfrak{G}_{d_t}(\Gamma) = \pi_*(\rho^*(\Gamma) \cdot \mathfrak{G}_{d_1} \dots \mathfrak{G}_{d_t}) : A^*(\text{Hilb}) \rightarrow A^*(\text{Hilb})$$

(where $\rho : \text{Hilb} \times S^t \rightarrow S^t$ is the second projection). By a push-pull argument, the expression above is the operator of multiplication by the universal class (2.9):

$$\mathfrak{G}_{d_1} \dots \mathfrak{G}_{d_t}(\Gamma) = \text{mult}_{\text{univ}_{d_1, \dots, d_t}}(\Gamma)$$

which explains our interest in the operators (4.3).

4.3. Consider the following operators, defined by [19] for all $n \in \mathbb{Z}$ and $d \in \mathbb{N} \sqcup 0$:

$$\mathfrak{F}_n^d : A^*(\text{Hilb}) \rightarrow A^*(\text{Hilb} \times S)$$

given by

$$(4.4) \quad \mathfrak{F}_n^d = d! \left(- \sum_{|\lambda|=n, l(\lambda)=d+1} \frac{\mathfrak{q}_\lambda}{\lambda!} \Big|_{\Delta} + \sum_{|\lambda|=n, l(\lambda)=d-1} \frac{s(\lambda) + n^2 - 2}{\lambda!} \rho^*(c) \mathfrak{q}_\lambda \Big|_{\Delta} \right),$$

where for any integer partition $\lambda = (\dots, (-2)^{m-2}, (-1)^{m-1}, 1^{m_1}, 2^{m_2}, \dots)$, we define

$$l(\lambda) = \sum_{i \in \mathbb{Z} \setminus 0} m_i, \quad |\lambda| = \sum_{i \in \mathbb{Z} \setminus 0} i m_i, \quad s(\lambda) = \sum_{i \in \mathbb{Z} \setminus 0} i^2 m_i, \quad \lambda! = \prod_{i \in \mathbb{Z} \setminus 0} m_i!$$

$$\mathfrak{q}_\lambda = \dots \mathfrak{q}_2^{m_2} \mathfrak{q}_1^{m_1} \mathfrak{q}_{-1}^{m-1} \mathfrak{q}_{-2}^{m-2} \dots : A^*(\text{Hilb}) \rightarrow A^*(\text{Hilb} \times S^{l(\lambda)})$$

and $|_{\Delta}$ denotes the restriction to the small diagonal $A^*(\text{Hilb} \times S^{l(\lambda)}) \rightarrow A^*(\text{Hilb} \times S)$. For any $\gamma \in A^*(S)$, we may consider the operator

$$\mathfrak{F}_n^d(\gamma) = \pi_*(\rho^*(\gamma) \cdot \mathfrak{F}_n^d) : A^*(\text{Hilb}) \rightarrow A^*(\text{Hilb})$$

and then formula (4.4) yields the following:

$$(4.5) \quad \mathfrak{F}_n^d(\gamma) = d! \left(- \sum_{|\lambda|=n, l(\lambda)=d+1} \frac{1}{\lambda!} \cdot \mathfrak{q}_\lambda(\Delta_{1\dots d+1} \gamma_1) + \sum_{|\lambda|=n, l(\lambda)=d-1} \frac{s(\lambda) + n^2 - 2}{\lambda!} \cdot \mathfrak{q}_\lambda(\Delta_{1\dots d-1} \gamma_1 c_1) \right),$$

where $\Delta_{1\dots d}$ denotes the small diagonal in S^d .

4.4. The following result is proved just like its cohomological counterpart in [19, Theorem 5.5] (the only input the computation needs is relation (2.14), which takes the same form in cohomology as in Chow).

Theorem 4.1. *For all $n, n' \in \mathbb{Z}$ and $d + d' \geq 3$ (or $d + d' = 2$ but $n + n' \neq 0$), we have*

$$(4.6) \quad [\mathfrak{F}_n^d, \mathfrak{F}_{n'}^{d'}] = (dn' - d'n)\Delta_*(\mathfrak{F}_{n+n'}^{d+d'-1}) + 2\Omega_{n,n'}^{d,d'}\Delta_*(\rho^*(c) \cdot \mathfrak{F}_{n+n'}^{d+d'-3})$$

as operators $A^*(\text{Hilb}) \rightarrow A^*(\text{Hilb} \times S \times S)$, where $\Delta : S \hookrightarrow S \times S$ is the diagonal, $\Omega_{n,n'}^{d,d'}$ are certain integers, and $\rho : \text{Hilb} \times S \rightarrow S$ is the second projection.

The precise formula for the numbers $\Omega_{n,n'}^{d,d'}$ can be found in [19, relation (5.2)] (note that one must replace $n, n' \leftrightarrow -n, -n'$ to match our notation with [19]), but we will only need the following particularly simple cases of formula (4.6):

$$(4.7) \quad [\mathfrak{q}_n, \mathfrak{F}_0^d] = dn\Delta_*(\mathfrak{F}_n^{d-1}),$$

$$(4.8) \quad [\mathfrak{L}_n, \mathfrak{F}_0^d] = dn\Delta_*(\mathfrak{F}_n^d) + 2d(d-1)n(n^2-1)\Delta_*(\rho^*(c) \cdot \mathfrak{F}_n^{d-2}),$$

where we note that $\mathfrak{F}_n^0 = -\mathfrak{q}_n$, while $\mathfrak{F}_n^1 = -\mathfrak{L}_n$ with

$$(4.9) \quad \mathfrak{L}_n = \frac{1}{2} \sum_{i,j \in \mathbb{Z}, i+j=n} : \mathfrak{q}_i \mathfrak{q}_j \Big|_{\Delta} :$$

and $:-$ denotes the normal ordered product (3.2).

Theorem 4.2 ([19, Theorem 4.6] in cohomology, [23, Theorem 1.7] in Chow). *For any $d \in \mathbb{N}$, we have*

$$\mathfrak{F}_0^d = d!(\mathfrak{G}_{d+1} + 2\rho^*(c) \cdot \mathfrak{G}_{d-1})$$

as operators $A^*(\text{Hilb}) \rightarrow A^*(\text{Hilb} \times S)$, where $\rho : \text{Hilb} \times S \rightarrow S$ is the second projection. Equivalently, we may write the formula above as

$$(4.10) \quad \mathfrak{F}_0^d(\gamma) = d!(\mathfrak{G}_{d+1}(\gamma) + 2\mathfrak{G}_{d-1}(\gamma c))$$

as operators $A^*(\text{Hilb}) \rightarrow A^*(\text{Hilb})$ indexed by $\gamma \in A^*(S)$.

Because $c^2 = 0$, inverting formula (4.10) gives

$$\mathfrak{G}_d = \frac{\mathfrak{F}_0^{d-1}}{(d-1)!} - \frac{2\rho^*(c) \cdot \mathfrak{F}_0^{d-3}}{(d-3)!}$$

or equivalently as

$$(4.11) \quad \mathfrak{G}_d(\gamma) = \frac{\mathfrak{F}_0^{d-1}(\gamma)}{(d-1)!} - \frac{2\mathfrak{F}_0^{d-3}(\gamma c)}{(d-3)!}.$$

4.5. Let us now recall the operators

$$(4.12) \quad h, h_{\alpha\beta}, h_{\alpha\delta} : A^*(\text{Hilb}) \rightarrow A^*(\text{Hilb})$$

defined in formula (3.4), (3.5), and (3.6) for all $\alpha, \beta \in A^1(S) \subset A^1(\text{Hilb})$. In order to prove Theorems 1.4 and 1.6, we need to compute the commutators of these operators with the operators (4.3) of multiplication by universal classes.

Proposition 4.3. *For any $d \geq 2$ and $\alpha, \beta \in A^1(S) \subset A^1(\text{Hilb})$, we have*

$$(4.13) \quad [h, \mathfrak{G}_d(\gamma)] = \mathfrak{G}_d \left((d-1)\gamma + \int_{\bullet} \gamma_{\bullet} (c - c_{\bullet}) \right),$$

$$(4.14) \quad [h_{\alpha\beta}, \mathfrak{G}_d(\gamma)] = \mathfrak{G}_d \left(\int_{\bullet} \gamma_{\bullet} (\alpha\beta_{\bullet} - \alpha_{\bullet}\beta) \right)$$

as operators $A^*(\text{Hilb}) \rightarrow A^*(\text{Hilb})$ indexed by $\gamma \in A^*(S)$.

Formula (4.13) is the main thing we need to prove Theorem 1.4. However, to prove Theorem 1.6, we will also need the following more complicated version of the formulas above.

Proposition 4.4. *For any $d \geq 2$ and $\alpha \in A^1(S) \subset A^1(\text{Hilb})$, we have*

$$(4.15) \quad [h_{\alpha\delta}, \mathfrak{G}_d(\gamma)] = -\mathfrak{G}_2(\alpha)\mathfrak{G}_{d-1}(\gamma) - \mathfrak{G}_2(1)\mathfrak{G}_{d-1}(\gamma\alpha) \\ - \mathfrak{G}_{d+1} \left(\alpha \int_{\bullet} \gamma_{\bullet} + \int_{\bullet} \gamma_{\bullet} \alpha_{\bullet} \right) + 2\mathfrak{G}_{d-1} \left(\alpha \int_{\bullet} \gamma_{\bullet} c_{\bullet} \right)$$

as operators $A^*(\text{Hilb}) \rightarrow A^*(\text{Hilb})$ indexed by $\gamma \in A^*(S)$.

We separate the relations above into two different propositions, because the proof of the latter will be significantly more involved than the former. But before we dive into the proof, let us observe that according to our conventions, we are actually proving (4.13), (4.14), and (4.15) as the following identities:

$$(4.16) \quad (h \times \text{Id}_S) \circ \mathfrak{G}_d - \mathfrak{G}_d \circ h = (d-1)\mathfrak{G}_d + \pi_{2*}((c_1 - c_2) \cdot \pi_1^*(\mathfrak{G}_d)),$$

$$(4.17) \quad (h_{\alpha\beta} \times \text{Id}_S) \circ \mathfrak{G}_d - \mathfrak{G}_d \circ h_{\alpha\beta} = \pi_{2*}((\alpha_1\beta_2 - \alpha_2\beta_1) \cdot \pi_1^*(\mathfrak{G}_d))$$

and

$$(4.18) \quad (h_{\alpha\delta} \times \text{Id}_S) \circ \mathfrak{G}_d - \mathfrak{G}_d \circ h_{\alpha\delta} \\ = \pi_{2*} \left[-(\alpha_1 + \alpha_2) \cdot \pi_1^*(\mathfrak{G}_2)\pi_2^*(\mathfrak{G}_{d-1}) - (\alpha_1 + \alpha_2) \cdot \pi_1^*(\mathfrak{G}_{d+1}) \right. \\ \left. + 2\alpha_1 c_2 \cdot \pi_1^*(\mathfrak{G}_{d-1}) \right]$$

of operators $A^*(\text{Hilb}) \rightarrow A^*(\text{Hilb} \times S)$, where $\pi_i : \text{Hilb} \times S \times S \rightarrow \text{Hilb} \times S$ denotes the identity on Hilb times the projection onto the i -th factor of S . The reason why we prefer the language of (4.13), (4.14), and (4.15) over (4.16), (4.17), and (4.18) is simply to keep the explanation legible.

4.6. The reader who is willing to accept Propositions 4.3 and 4.4, and wishes to see how they lead to Theorems 1.4 and 1.6, may skip to Section 4.8.

Proof of Proposition 4.3. We will start with (4.13). From a straightforward calculation (see [29, Lemma 3.4]), one obtains the commutation relations

$$(4.19) \quad [h, \mathfrak{q}_{\lambda_1} \cdots \mathfrak{q}_{\lambda_k}(\Phi)] = \mathfrak{q}_{\lambda_1} \cdots \mathfrak{q}_{\lambda_k}(\bar{\Phi})$$

for all $\Phi \in A^*(S^k)$, where we write

$$(4.20) \quad \bar{\Phi} = \sum_{i=1}^k \int_{\bullet} \underbrace{\Phi_{1\dots i-1, \bullet, i+1\dots k}(c_i - c_{\bullet})}_{\text{this class lies in } A^*(S^k \times S)}$$

with the last factor in $S^k \times S$ represented by the index \bullet . We have

$$(4.21) \quad [h, \mathfrak{S}_0^d(\gamma)] \stackrel{(4.5)}{=} d! \left(- \sum_{|\lambda|=0, l(\lambda)=d+1} \frac{1}{\lambda!} \cdot [h, \mathfrak{q}_\lambda(\Delta_{1\dots d+1}\gamma_1)] \right. \\ \left. + \sum_{|\lambda|=0, l(\lambda)=d-1} \frac{s(\lambda)-2}{\lambda!} \cdot [h, \mathfrak{q}_\lambda(\Delta_{1\dots d-1}\gamma_1 c_1)] \right) \\ \stackrel{(4.19)}{=} d! \left(- \sum_{|\lambda|=0, l(\lambda)=d+1} \frac{1}{\lambda!} \cdot \mathfrak{q}_\lambda(\overline{\Delta_{1\dots d+1}\gamma_1}) \right. \\ \left. + \sum_{|\lambda|=0, l(\lambda)=d-1} \frac{s(\lambda)-2}{\lambda!} \cdot \mathfrak{q}_\lambda(\overline{\Delta_{1\dots d-1}\gamma_1 c_1}) \right).$$

To evaluate the expression above, we will need the result below:

Claim 4.5. For any $k > 0$, $l \geq 0$ and any $\gamma \in A^*(S \times S^l)$, we have

$$(4.22) \quad \overline{\Delta_{1\dots k}\gamma_1} = \Delta_{1\dots k} \left[(k-1)\gamma_1 + \int_* \gamma_*(c_1 - c_*) \right].$$

When writing γ_* , the index $*$ refers to the first factor of $\gamma \in A^*(S \times S^l)$. The other l factors of γ are not involved in the formula above, as the bar notation is defined as in (4.20) with respect to the indices $1, \dots, k$ only.

Proof. In the sequel, we let \bullet and $*$ denote two different copies of the surface S which will be integrated out. Using (2.7), the left-hand side of (4.22) equals

$$\sum_{i=1}^k \int_{\bullet} \Delta_{1\dots i-1, \bullet, i+1\dots k} \gamma_{\bullet}(c_i - c_{\bullet}) \\ = \left[\sum_{i=1}^k \sum_{j \in \{1, \dots, k, \bullet\} - \{i\}} \int_{\bullet} \gamma_j \prod_{x \neq i, j} c_x (c_i - c_{\bullet}) \right] \\ + \left[\sum_{i=1}^k \int_{\bullet} \left(\Delta_{1\dots i-1, \bullet, i+1\dots k} - \sum_{j \in \{1, \dots, k, \bullet\} - \{i\}} \prod_{x \neq i, j} c_x \right) (c_i - c_{\bullet}) \int_* \gamma_* c_* \right] \\ - \left[\sum_{i=1}^k (k-1) \int_{\bullet} c_1 \dots c_{i-1} c_{\bullet} c_{i+1} \dots c_k (c_i - c_{\bullet}) \int_* \gamma_* \right].$$

A straightforward calculation using formula (2.5) shows that the three square brackets above are equal to the corresponding three square brackets below (one needs to integrate out the factor denoted by \bullet):

$$(4.23) \quad \left[(k-1) \sum_{i=1}^k \gamma_i \prod_{j \neq i} c_j + k c_1 \dots c_k \int_* \gamma_* - \sum_{i=1}^k \prod_{j \neq i} c_j \int_* \gamma_* c_* \right] \\ + \left[\sum_{i=1}^k \left(\Delta_{1\dots \hat{i} \dots k} c_i - (k-1) \prod_{j \neq i} c_j \right) \int_* \gamma_* c_* \right] \\ - \left[k(k-1) c_1 \dots c_k \int_* \gamma_* \right].$$

Using the following corollary of (2.4):

$$\sum_{i=1}^k \Delta_{1 \dots \hat{i} \dots k} c_i = (k-2) \Delta_{1 \dots k} + \sum_{i=1}^k c_1 \dots c_{i-1} c_{i+1} \dots c_k,$$

we may rearrange (4.23) as

$$\begin{aligned} & (k-1) \sum_{i=1}^k \gamma_i \prod_{j \neq i} c_j - [k(k-2)c_1 \dots c_k] \int_* \gamma_* \\ & + \left[(k-2) \Delta_{1 \dots k} - (k-1) \sum_{i=1}^k \prod_{j \neq i} c_j \right] \int_* \gamma_* c_*. \end{aligned}$$

Using formulas (2.5) and (2.7), one recognizes that the expression above equals

$$\Delta_{1 \dots k} \left[(k-1) \gamma_1 + \int_* \gamma_* (c_1 - c_*) \right]$$

thus establishing formula (4.22). \square

Relation (4.21) together with the following immediate consequence of (2.5):

$$-2\gamma c + \int_* \gamma_* c_* (c - c_*) = \int_* \gamma_* c (c - c_*)$$

implies that

$$(4.24) \quad [h, \mathfrak{F}_0^d(\gamma)] = \mathfrak{F}_0^d \left(d\gamma + \int_* \gamma_* (c - c_*) \right).$$

Together with (4.11), formula (4.24) implies (4.13).

In order to prove (4.14), we will recycle the argument above. By analogy with (4.19), we have

$$[h_{\alpha\beta}, \mathfrak{q}_{\lambda_1} \dots \mathfrak{q}_{\lambda_k}(\Phi)] = \mathfrak{q}_{\lambda_1} \dots \mathfrak{q}_{\lambda_k}(\overline{\overline{\Phi}})$$

for all $\Phi \in A^*(S^k)$, where we write

$$(4.25) \quad \overline{\overline{\Phi}} = \sum_{i=1}^k \int_{\bullet} \underbrace{\Phi_{1 \dots i-1, \bullet, i+1 \dots k}}_{\text{this class lies in } A^*(S^k \times S)} (\alpha_i \beta_{\bullet} - \alpha_{\bullet} \beta_i),$$

where the last factor in $S^k \times S$ is the one represented by the index \bullet .

Claim 4.6. *For any $k > 0$, $l \geq 0$ and any $\gamma \in A^*(S \times S^l)$, we have*

$$(4.26) \quad \overline{\overline{\Delta_{1 \dots k} \gamma_1}} = \Delta_{1 \dots k} \int_* \gamma_* (\alpha_1 \beta_* - \alpha_* \beta_1).$$

When writing γ_* , the index $*$ refers to the first factor of $\gamma \in A^*(S \times S^l)$. The other l factors of γ are not involved in the formula above, as the double bar notation is defined as in (4.25) with respect to the indices $1, \dots, k$ only.

Proof. In the sequel, we let \bullet and $*$ denote two different copies of the surface S which will be integrated out. By definition, the left-hand side of (4.26) equals

$$\begin{aligned} & \sum_{i=1}^k \int_{\bullet} \Delta_{1\dots i-1, \bullet, i+1\dots k} \gamma_{\bullet} (\alpha_i \beta_{\bullet} - \alpha_{\bullet} \beta_i) \\ & \stackrel{(2.7)}{=} \left[\sum_{i=1}^k \sum_{j \in \{1, \dots, k, \bullet\} - \{i\}} \int_{\bullet} \gamma_j \prod_{x \neq i, j} c_x (\alpha_i \beta_{\bullet} - \alpha_{\bullet} \beta_i) \right] \\ & \quad + \left[\sum_{i=1}^k \int_{\bullet} \left(\Delta_{1\dots i-1, \bullet, i+1\dots k} - \sum_{j \in \{1, \dots, k, \bullet\} - \{i\}} \prod_{x \neq i, j} c_x \right) (\alpha_i \beta_{\bullet} - \alpha_{\bullet} \beta_i) \int_{*} c_{*} \gamma_{*} \right] \\ & \quad - \left[\sum_{i=1}^k (k-1) \int_{\bullet} c_1 \dots c_{i-1} c_{\bullet} c_{i+1} \dots c_k (\alpha_i \beta_{\bullet} - \alpha_{\bullet} \beta_i) \int_{*} \gamma_{*} \right]. \end{aligned}$$

One can apply (2.6) to compute the square brackets above (integrate out the factor of S denoted by \bullet), and obtain

$$= \left[\sum_{i=1}^k \prod_{j \neq i} c_j \left(\alpha_i \int_{*} \gamma_{*} \beta_{*} - \beta_i \int_{*} \gamma_{*} \alpha_{*} \right) \right] + [0] + [0].$$

Formula (2.2) shows that the right-hand side of the formula above is equal to

$$\Delta_{1\dots k} \int_{*} \gamma_{*} (\alpha_1 \beta_{*} - \alpha_{*} \beta_1)$$

which establishes (4.26). \square

The analogue of formula (4.21) holds with h replaced by $h_{\alpha\beta}$ and the bar replaced by a double bar, hence Claim 4.6 implies the following analogue of formula (4.24):

$$[h_{\alpha\beta}, \mathfrak{F}_0^d(\gamma)] = \mathfrak{F}_0^d \left(\int_{\bullet} \gamma_{\bullet} (\alpha \beta_{\bullet} - \alpha_{\bullet} \beta) \right).$$

Together with (4.11), this implies (4.14). \square

4.7. Proof of Proposition 4.4. Recall the definition of the operators of \mathfrak{L}_k in (4.9). Then formula (3.6) takes the form

$$\begin{aligned} (4.27) \quad h_{\alpha\delta} &= \sum_{k \neq 0} \frac{1}{k} : \mathfrak{L}_k \mathfrak{q}_{-k} (\alpha_1 + \alpha_2) : \\ &= \sum_{(x,y) \in \{(\alpha,1), (1,\alpha)\}} \sum_{k \neq 0} \frac{1}{k} : \mathfrak{L}_k(x) \mathfrak{q}_{-k}(y) :. \end{aligned}$$

We may invoke (4.7) and (4.8) to obtain

$$\begin{aligned} [h_{\alpha\delta}, \mathfrak{F}_0^d(\gamma)] &= d \sum_{k \neq 0} \sum_{(x,y) \in \{(\alpha,1), (1,\alpha)\}} \left[- : \mathfrak{L}_k(x) \mathfrak{F}_{-k}^{d-1}(\gamma\gamma) : + : \mathfrak{F}_k^d(x\gamma) \mathfrak{q}_{-k}(y) : \right. \\ & \quad \left. + 2(d-1)(k^2-1) : \mathfrak{F}_k^{d-2}(x\gamma) \mathfrak{q}_{-k}(y) : \right]. \end{aligned}$$

In the normal ordered products above, we put \mathfrak{q}_k and \mathfrak{L}_k to the left of the expression if $k > 0$ and to the right of the expression if $k < 0$. Since $\mathfrak{q}_0 = 0$ but $\mathfrak{L}_0 = -\mathfrak{J}_0^1$, we may rewrite the formula above as

$$(4.28) \quad [h_{\alpha\delta}, \mathfrak{J}_0^d(\gamma)] = -d \mathfrak{J}_0^1(\alpha) \mathfrak{J}_0^{d-1}(\gamma) - d \mathfrak{J}_0^1(1) \mathfrak{J}_0^{d-1}(\gamma\alpha) \\ + d \sum_{k \in \mathbb{Z}} \sum_{(x,y) \in \{(\alpha,1), (1,\alpha)\}} [-:\mathfrak{L}_k(y) \mathfrak{J}_{-k}^{d-1}(x\gamma): \\ + : \mathfrak{J}_k^d(x\gamma) \mathfrak{q}_{-k}(y): + 2(d-1)(k^2-1) : \mathfrak{J}_k^{d-2}(xc\gamma) \mathfrak{q}_{-k}(y):].$$

Let us compute the formulas on the second and third lines of the formula above.

Claim 4.7. *We have the following formulas:*

$$(4.29) \quad \sum_{k \in \mathbb{Z}} \sum_{|\lambda|=k, l(\lambda)=d+1} \frac{1}{\lambda!} : \mathfrak{q}_\lambda(\Delta_{1\dots d+1}(x\gamma)_1) \mathfrak{q}_{-k}(y): \\ = \sum_{\substack{|\mu|=0 \\ l(\mu)=d+2}} \frac{1}{\mu!} \cdot \mathfrak{q}_\mu \left(\sum_{i=1}^{d+2} \Delta_{1\dots \hat{i} \dots d+2}(x\gamma)_{\neq i} y_i \right)$$

and

$$(4.30) \quad \sum_{k \in \mathbb{Z}} \sum_{|\lambda|=k, l(\lambda)=d-1} \frac{s(\lambda) + k^2 - 2}{\lambda!} : \mathfrak{q}_\lambda(\Delta_{1\dots d-1}(x\gamma c)_1) \mathfrak{q}_{-k}(y): \\ = \sum_{|\mu|=0, l(\mu)=d} \frac{s(\mu) - 2}{\mu!} \mathfrak{q}_\mu \left(\sum_{i=1}^d \Delta_{1\dots \hat{i} \dots d}(x\gamma c)_{\neq i} y_i \right),$$

where $\Delta_{1\dots \hat{i} \dots d} x_{\neq i}$ refers to $\Delta_{1\dots \hat{i} \dots d} x_j$ for any $j \neq i$.

Proof. The right-hand side of (4.29) is equal to

$$\sum_{\mu=(\dots, (-2)^{m-2}, (-1)^{m-1}, 1^{m_1}, 2^{m_2}, \dots)} \frac{\dots \mathfrak{q}_{-k}^{m-k} \dots}{\dots m-k! \dots} \left(\sum_{i=1}^{l(\mu)} y_i \Delta_{\dots \hat{i} \dots} (x\gamma)_{\neq i} \right).$$

In each summand above, we can pick a copy of \mathfrak{q}_{-m_k} in $m-k$ ways for any k , and assign to that copy the insertion y_i , and to all other copies the insertion $\Delta_{\dots \hat{i} \dots} (x\gamma)_{\neq i}$. The corresponding sum will be term-wise equal to the left-hand side of (4.29). Formula (4.30) is proved analogously, so we leave it to the interested reader. \square

Claim 4.8. *We have the following formulas:*

$$(4.31) \quad \sum_{k \in \mathbb{Z}} \sum_{|\lambda|=-k, l(\lambda)=d} \frac{1}{\lambda!} : \mathfrak{L}_k(y) \mathfrak{q}_\lambda(\Delta_{1\dots d}(x\gamma)_1): \\ = \sum_{|\mu|=0, l(\mu)=d+2} \frac{1}{\mu!} \mathfrak{q}_\mu \left(\sum_{1 \leq i < j \leq d+2} \Delta_{1\dots \hat{i} \dots \hat{j} \dots d+2}(x\gamma)_{\neq i, j} \Delta_{ij} y_i \right) \\ - \sum_{|\mu|=0, l(\mu)=d} \frac{s(\mu)}{2\mu!} \mathfrak{q}_\mu(\Delta_{1\dots d}(xy\gamma)_1)$$

and

$$\begin{aligned}
(4.32) \quad & \sum_{k \in \mathbb{Z}} \sum_{|\lambda|=-k, l(\lambda)=d-2} \frac{s(\lambda) + k^2 - 2}{\lambda!} : \mathfrak{L}_k(y) \mathfrak{q}_\lambda(\Delta_{1\dots d-2}(xc\gamma)_1) : \\
& = \sum_{|\mu|=0, l(\mu)=d} \frac{s(\mu) - 2}{\mu!} \mathfrak{q}_\mu \left(\sum_{1 \leq i < j \leq d} \Delta_{1\dots \hat{i} \dots \hat{j} \dots d}(xc\gamma)_{\neq i, j} \Delta_{ij} y_i \right) \\
& \quad + \sum_{i, j \in \mathbb{Z}} \sum_{|\lambda|=-i-j, l(\lambda)=d-2} \frac{ij}{\lambda!} : \mathfrak{q}_i \mathfrak{q}_j(\Delta_{12} y_1) \mathfrak{q}_\lambda(\Delta_{1\dots d-2}(xc\gamma)_1) :.
\end{aligned}$$

Proof. As in the proof of (4.29), the terms on the first line of (4.31) are in one-to-one correspondence with the terms on the second line. However, while the latter are normally ordered by definition, the former are not always normally ordered, due to the presence of the following terms:

$$(4.33) \quad \mathfrak{q}_k \mathfrak{q}_{-l}(\Delta_{12} y_1) \mathfrak{q}_\lambda(\Delta_{1\dots d}(x\gamma)_1),$$

$$(4.34) \quad \mathfrak{q}_\lambda(\Delta_{1\dots d}(x\gamma)_1) \mathfrak{q}_l \mathfrak{q}_{-k}(\Delta_{12} y_1)$$

for all $k \geq l \geq 0$ (if $k = l$, the corresponding product appears in both (4.33) and (4.34), and we weigh it with weight $\frac{1}{2}$ in both of these formulas). Therefore, the difference between the first and second lines of (4.31) is equal to the work necessary in normally ordering the expressions (4.33) and (4.34), and we must identify the contribution of these with the expression on the third line of (4.31). For fixed k , this contribution is

$$\begin{aligned}
& - \left(1 + 2 + \dots + k - 1 + \frac{k}{2} \right) \mathfrak{q}_k \mathfrak{q}_{\lambda'}(\Delta_{1\dots d}(xy\gamma)_1) \quad \text{in the case (4.33),} \\
& - \left(1 + 2 + \dots + k - 1 + \frac{k}{2} \right) \mathfrak{q}_{\lambda''} \mathfrak{q}_{-k}(\Delta_{1\dots d}(xy\gamma)_1) \quad \text{in the case (4.34),}
\end{aligned}$$

where λ' (respectively λ'') denotes λ without any one factor \mathfrak{q}_l with l positive (respectively negative). As we sum over all partitions λ and over all ways to remove any one factor \mathfrak{q}_l from them, we are left with

$$- \sum_{k \in \mathbb{Z}} \sum_{|\lambda|=-k, l(\lambda)=d-1} \frac{k^2}{2\lambda!} : \mathfrak{q}_k \mathfrak{q}_\lambda(\Delta_{1\dots d}(xy\gamma)_1) :,$$

which is precisely the third line of (4.31). Formula (4.32) is proved analogously, only that we do not have to worry about the commutators that arose in the preceding paragraph, because $xy\gamma = 0$ for any $(x, y) \in \{(\alpha, 1), (1, \alpha)\}$. The expression on the last line of (4.32) simply arises as the difference $k^2 - i^2 - j^2 = s(\lambda) + k^2 - s(\lambda \sqcup \{i, j\})$ in the notation thereof. \square

For every $k \in \mathbb{N}$ consider the cycles in S^k defined by

$$\begin{aligned}
A_k(\gamma) & = \sum_{(x, y) \in \{(\alpha, 1), (1, \alpha)\}} \sum_{1 \leq i < j \leq k} \Delta_{1\dots \hat{i} \dots \hat{j} \dots k}(x\gamma)_{\neq i, j} \Delta_{ij} y_i, \\
B_k(\gamma) & = \sum_{(x, y) \in \{(\alpha, 1), (1, \alpha)\}} \sum_{i=1}^k \Delta_{1\dots \hat{i} \dots k}(x\gamma)_{\neq i} y_i.
\end{aligned}$$

Then using Claims 4.7 and 4.8 and (4.5), we may rewrite formula (4.28) as

$$\begin{aligned}
 (4.35) \quad [h_{\alpha\delta}, \mathfrak{F}_0^d(\gamma)] &= -d \mathfrak{F}_0^1(\alpha) \mathfrak{F}_0^{d-1}(\gamma) - d \mathfrak{F}_0^1(1) \mathfrak{F}_0^{d-1}(\gamma\alpha) + d! \left[\sum_{l(\mu)=d+2} \frac{1}{\mu!} \mathfrak{q}_\mu(A_{d+2}(\gamma)) \right. \\
 &\quad - \sum_{(x,y)} \sum_{l(\mu)=d} \frac{s(\mu)}{2\mu!} \mathfrak{q}_\mu(\Delta_{1\dots d}(xy\gamma)_1) - \sum_{l(\mu)=d} \frac{s(\mu)-2}{\mu!} \mathfrak{q}_\mu(A_d(c\gamma)) \\
 &\quad - \sum_{(x,y)} \sum_{i,j \in \mathbb{Z}} \sum_{|\lambda|=-i-j, l(\lambda)=d-2} \frac{ij}{\lambda!} : \mathfrak{q}_i \mathfrak{q}_j(\Delta_{12}y_1) \mathfrak{q}_\lambda(\Delta_{1\dots d-2}(xc\gamma)_1) : \\
 &\quad - d \sum_{l(\mu)=d+2} \frac{1}{\mu!} \mathfrak{q}_\mu(B_{d+2}(\gamma)) + d \sum_{l(\mu)=d} \frac{s(\mu)-2}{\mu!} \mathfrak{q}_\mu(B_d(c\gamma)) \\
 &\quad \left. - \sum_{(x,y)} \sum_{k \in \mathbb{Z}} \sum_{|\lambda|=k, l(\lambda)=d-1} \frac{2(k^2-1)}{\lambda!} : \mathfrak{q}_\lambda(\Delta_{1\dots d-1}(xc\gamma)_1) \mathfrak{q}_{-k}(y) : \right],
 \end{aligned}$$

where above and hereafter, all the partitions denoted by μ will have $|\mu| = 0$ and (x, y) runs over $\{(\alpha, 1), (1, \alpha)\}$.

Claim 4.9. *The sum of the third and fifth lines of (4.35) equals*

$$(4.36) \quad 2 \sum_{l(\mu)=d} \frac{1}{\mu!} \mathfrak{q}_\mu \left(\sum_{i=1}^d \Delta_{1\dots \hat{i} \dots d}(c\gamma)_{\neq i} \alpha_i \right).$$

Proof. Because $\alpha c = 0$, only the $(x, y) = (1, \alpha)$ has a non-zero contribution to the third and fifth lines of (4.35), which means that their sum equals (using (2.1) and (2.2))

$$\begin{aligned}
 (4.37) \quad & - \sum_{k \in \mathbb{Z}} \sum_{|\lambda|=k, l(\lambda)=d-1} \frac{2(k^2-1)}{\lambda!} : \mathfrak{q}_\lambda(\gamma_1 c_1 \dots c_{d-1}) \mathfrak{q}_{-k}(\alpha) : \\
 & - \sum_{i,j \in \mathbb{Z}} \sum_{|\lambda|=-i-j, l(\lambda)=d-2} \frac{ij}{\lambda!} : \mathfrak{q}_i \mathfrak{q}_j(\alpha_1 c_2 + c_1 \alpha_2) \mathfrak{q}_\lambda(\gamma_1 c_1 \dots c_{d-2}) :.
 \end{aligned}$$

Because $\alpha c = 0$, all the \mathfrak{q}_i commute in the formula above, hence the second line of (4.37) equals

$$\underline{-2 \sum_{i \in \mathbb{Z}} : i \mathfrak{q}_i(\alpha) \sum_{j \in \mathbb{Z}} \sum_{|\lambda|=-i-j, l(\lambda)=d-2} \frac{j}{\lambda!} \mathfrak{q}_j(c) \mathfrak{q}_\lambda(\gamma_1 c_1 \dots c_{d-2}) :.}$$

The underlined sum is equal to

$$\sum_{|\lambda|=-i, l(\lambda)=d-1} \frac{|\lambda|}{\lambda!} \mathfrak{q}_\lambda(\gamma_1 c_1 \dots c_{d-1}).$$

Plugging this fact into (4.37) leads to

$$2 \sum_{k \in \mathbb{Z}} \sum_{|\lambda|=k, l(\lambda)=d-1} \frac{1}{\lambda!} : \mathfrak{q}_\lambda(\gamma_1 c_1 \dots c_{d-1}) \mathfrak{q}_{-k}(\alpha) :$$

which is equal to (4.36) by a straightforward rearranging of terms (akin to the one we performed in Claim 4.7). \square

Using Claim 4.9, and after reordering terms, we may rewrite (4.35) as

$$(4.38) \quad [h_{\alpha\delta}, \mathfrak{F}_0^d(\gamma)] = -d \mathfrak{F}_0^1(\alpha) \mathfrak{F}_0^{d-1}(\gamma) - d \mathfrak{F}_0^1(1) \mathfrak{F}_0^{d-1}(\gamma\alpha) \\ + d! \left[\sum_{l(\mu)=d+2} \frac{1}{\mu!} \mathfrak{q}_\mu(A_{d+2}(\gamma) - dB_{d+2}(\gamma)) \right. \\ + 2 \sum_{l(\mu)=d} \frac{1}{\mu!} \mathfrak{q}_\mu(A_d(\gamma c) - (d-1)B_d(\gamma c)) \\ + \sum_{l(\mu)=d} \frac{s(\mu)}{\mu!} \mathfrak{q}_\mu \left(dB_d(\gamma c) - A_d(\gamma c) \right. \\ \left. \left. - \Delta_{1\dots d} \left(\alpha_1 \int_{\bullet} \gamma_{\bullet} c_{\bullet} + c_1 \int_{\bullet} \gamma_{\bullet} \alpha_{\bullet} \right) \right) \right],$$

where in the last term, we used (2.6).

Lemma 4.10. *We have*

$$A_k(\gamma) - (k-2)B_k(\gamma) = \Delta_{1\dots k} \left(\alpha_1 \int_{\bullet} \gamma_{\bullet} + \int_{\bullet} \alpha_{\bullet} \gamma_{\bullet} \right)$$

and

$$A_k(\gamma c) = (k-1)\Delta_{1\dots k}\alpha_1 \int_{\bullet} \gamma_{\bullet} c_{\bullet}, \quad B_k(\gamma c) = \Delta_{1\dots k}\alpha_1 \int_{\bullet} \gamma_{\bullet} c_{\bullet}.$$

Proof. The equations in the second line follow immediately from (2.1). The first line follows from (2.4) and the following claim. \square

Claim 4.11. *For any $\alpha \in A^1(S)$ and any $\gamma \in A^*(S \times S^l)$, we have the following:*

$$(4.39) \quad A_k(\gamma) = (k-2) \sum_{i \neq j} \alpha_i \gamma_j \prod_{s \neq i, j} c_s - (k-1)(k-3) \left(\sum_i \alpha_i \prod_{j \neq i} c_j \right) \left(\int_{\bullet} \gamma_{\bullet} \right) \\ + \left(\sum_{i < j} \Delta_{ij} \prod_{s \neq i, j} c_s \right) \left(\int_{\bullet} \gamma_{\bullet} \alpha_{\bullet} \right) + (k-2) \left(\sum_{i \neq s < t \neq i} \alpha_i \Delta_{st} \prod_{j \neq i, s, t} c_j \right. \\ \left. - (k-3) \alpha_i \sum_{j \neq i} \prod_{s \neq i, j} c_s \right) \left(\int_{\bullet} \gamma_{\bullet} c_{\bullet} \right)$$

and

$$(4.40) \quad B_k(\gamma) = \sum_{i \neq j} \alpha_i \gamma_j \prod_{s \neq i, j} c_s - (k-2) \left(\sum_i \alpha_i \prod_{j \neq i} c_j \right) \left(\int_{\bullet} \gamma_{\bullet} \right) \\ + \left(\sum_i \prod_{j \neq i} c_j \right) \left(\int_{\bullet} \gamma_{\bullet} \alpha_{\bullet} \right) + \left(\sum_{i \neq s < t \neq i} \alpha_i \Delta_{st} \prod_{j \neq i, s, t} c_j \right. \\ \left. - (k-3) \alpha_i \sum_{j \neq i} \prod_{s \neq i, j} c_s \right) \left(\int_{\bullet} \gamma_{\bullet} c_{\bullet} \right).$$

Proof. Let us prove (4.40) and leave the analogous formula (4.39) as an exercise to the interested reader. Formulas (2.4) and (2.7) imply

$$\begin{aligned} \Delta_{\dots\hat{i}\dots}(x\gamma)_{\neq i} y_i &= \sum_{i \neq j} \gamma_j x_j y_i \prod_{s \neq i, j} c_s + \left(\sum_{i \neq s < t \neq i} \Delta_{st} y_i \prod_{j \neq i, s, t} c_j \right. \\ &\quad \left. - (k-2) y_i \sum_{j \neq i} \prod_{s \neq i, j} c_s \right) \left(\int_{\bullet} \gamma_{\bullet} x_{\bullet} c_{\bullet} \right) \\ &\quad - (k-2) y_i \prod_{j \neq i} c_j \left(\int_{\bullet} \gamma_{\bullet} x_{\bullet} \right). \end{aligned}$$

If we sum over $i \in \{1, \dots, k\}$ and over $(x, y) \in \{(\alpha, 1), (1, \alpha)\}$, we obtain (4.40) (note that in the $(x, y) = (\alpha, 1)$ case of the first term in the right-hand side, we need to use formula (2.6) to calculate $\gamma_j \alpha_j$). \square

With Lemma 4.10 in mind, (4.38) reads

$$\begin{aligned} [h_{\alpha\delta}, \mathfrak{F}_0^d(\gamma)] &= -d \mathfrak{F}_0^1(\alpha) \mathfrak{F}_0^{d-1}(\gamma) - d \mathfrak{F}_0^1(1) \mathfrak{F}_0^{d-1}(\gamma\alpha) \\ &\quad + d! \left[\sum_{l(\mu)=d+2} \frac{1}{\mu!} \mathfrak{q}_{\mu} \left(\Delta_{1\dots d+2} \left(\alpha_1 \int_{\bullet} \gamma_{\bullet} + \int_{\bullet} \gamma_{\bullet} \alpha_{\bullet} \right) \right) \right. \\ &\quad \left. - \sum_{l(\mu)=d} \frac{s(\mu)}{\mu!} \mathfrak{q}_{\mu} \left(\Delta_{1\dots d} c_1 \int_{\bullet} \gamma_{\bullet} \alpha_{\bullet} \right) \right]. \end{aligned}$$

By (4.4), the second and third lines of the expression above equal

$$d! \left[-\frac{1}{(d+1)!} \mathfrak{F}_0^{d+1} \left(\alpha \int_{\bullet} \gamma_{\bullet} + \int_{\bullet} \gamma_{\bullet} \alpha_{\bullet} \right) + \frac{2}{(d-1)!} \mathfrak{F}_0^{d-1} \left(c \int_{\bullet} \gamma_{\bullet} \alpha_{\bullet} \right) \right].$$

If we convert the operators \mathfrak{F} to operators \mathfrak{G} in the formula above using (4.10), we obtain formula (4.15). \square

4.8. Before we prove Theorems 1.4 and 1.6, let us compute how the operators (4.12) act on the fundamental class.

Lemma 4.12. *If $1_n \in A^*(\text{Hilb}_n)$ denotes the fundamental class, then $h(1_n) = -n$ and $h_{\alpha\beta}(1_n) = h_{\alpha\delta}(1_n) = 0$ for all $\alpha, \beta \in A^1(S) \subset A^1(X)$.*

Proof. It is wellknown that

$$1_n = \frac{1}{n!} \mathfrak{q}_1(1)^n(1_0).$$

Because the only operator \mathfrak{q}_k which fails to commute with \mathfrak{q}_1 is \mathfrak{q}_{-1} , formula (3.4) implies that

$$\begin{aligned} h(1_n) &= h \left(\frac{1}{n!} \mathfrak{q}_1(1)^n(1_0) \right) = \left[h, \frac{1}{n!} \mathfrak{q}_1(1)^n \right] (1_0) \\ &= \left[\mathfrak{q}_1(1) \mathfrak{q}_{-1}(c) - \mathfrak{q}_1(c) \mathfrak{q}_{-1}(1), \frac{1}{n!} \mathfrak{q}_1(1)^n \right] (1_0) \\ &= \sum_{i=1}^n \frac{1}{n!} \mathfrak{q}_1(1)^{i-1} \cdot \mathfrak{q}_1(1) [\mathfrak{q}_{-1}(c), \mathfrak{q}_1(1)] \cdot \mathfrak{q}_{-1}(1)^{n-i} \cdot (1_0) = -n 1_n, \end{aligned}$$

where the fact that $[\mathfrak{q}_{-1}(c), \mathfrak{q}_1(1)] = -1$ is a consequence of formula (2.15). The fact that $[\mathfrak{q}_{-1}(\alpha), \mathfrak{q}_1(1)] = 0$ for any $\alpha \in A^1(S)$ means that the analogous computation implies that $h_{\alpha\beta}(1_n) = 0$. Similarly, let us use formula (4.27) to compute

$$(4.41) \quad \begin{aligned} h_{a\delta}(1_n) &= \frac{1}{n!} [h_{a\delta}, \mathfrak{q}_1(1)^n](1_0) \\ &= \frac{2}{n!} \sum_{k=1}^{\infty} \frac{1}{k} [\mathfrak{L}_k \mathfrak{q}_{-k}(\alpha_1 + \alpha_2) - \mathfrak{q}_k \mathfrak{L}_{-k}(\alpha_1 + \alpha_2), \mathfrak{q}_1(1)^n](1_0). \end{aligned}$$

As a consequence of formula (2.15), $\mathfrak{q}_{-k}(\alpha)$ and $\mathfrak{q}_{-k}(1)$ commute with $\mathfrak{q}_1(1)$. Meanwhile, the well-known Virasoro algebra relation [18] reads $[\mathfrak{L}_k(\gamma), \mathfrak{q}_1(1)] = -\mathfrak{q}_{k+1}(\gamma)$ for all $k \in \mathbb{Z}$ and $\gamma \in A^*(S)$. Therefore, all the commutators vanish in (4.41) (or more precisely, they all have an annihilating operator \mathfrak{q}_{-k} with $k \geq 0$ on the very right, and therefore act by 0 on 1_0) hence $h_{a\delta}(1_n) = 0$. \square

4.9. By iterating relation (4.13) t times, we infer the following formula for all integers $d_1, \dots, d_t \geq 2$ and all $\Gamma \in A^*(S^t)$:

$$[h, \mathfrak{G}_{d_1} \dots \mathfrak{G}_{d_t}(\Gamma)] = \mathfrak{G}_{d_1} \dots \mathfrak{G}_{d_t} \left(\sum_{i=1}^t (d_i - 1)\Gamma + \int_{\bullet} \sum_{i=1}^t \Gamma_{1\dots i-1, \bullet, i+1\dots t} (c_i - c_{\bullet}) \right)$$

(note that we are actually using formula (4.16) in order to conclude the aforementioned result). Since $\mathfrak{G}_{d_1} \dots \mathfrak{G}_{d_t}(\Gamma)$ is the operator of multiplication by $\text{univ}_{d_1, \dots, d_t}(\Gamma)$, we conclude that

$$(4.42) \quad [h, \text{mult}_{\text{univ}_{d_1, \dots, d_t}(\Gamma)}] = \text{mult}_{\text{univ}_{d_1, \dots, d_t}(\Gamma')},$$

where $\Gamma' = \sum_{i=1}^t (d_i - 1)\Gamma + \int_{\bullet} \sum_{i=1}^t \Gamma_{1\dots i-1, \bullet, i+1\dots t} (c_i - c_{\bullet})$. We are now ready to prove Theorem 1.4, and follow the strategy outlined in Section 4.1.

Proof of Theorem 1.4. Let us first prove part (i). By applying (4.42) to the fundamental class $1_n \in A^*(\text{Hilb}_n)$, we obtain

$$h(\text{univ}_{d_1, \dots, d_t}(\Gamma)) - \text{univ}_{d_1, \dots, d_t}(\Gamma) \cdot h(1_n) = \text{univ}_{d_1, \dots, d_t}(\Gamma').$$

As a consequence of Lemma 4.12, we obtain

$$(4.43) \quad \begin{aligned} \text{univ}_{d_1, \dots, d_t}(\Gamma') &= h(\text{univ}_{d_1, \dots, d_t}(\Gamma)) + n \cdot \text{univ}_{d_1, \dots, d_t}(\Gamma) \\ &= \widetilde{h}(\text{univ}_{d_1, \dots, d_t}(\Gamma)) \end{aligned}$$

on $A^*(\text{Hilb}_n)$. Therefore, relation (4.42) reads

$$(4.44) \quad [\widetilde{h}, \text{mult}_{\text{univ}_{d_1, \dots, d_t}(\Gamma)}] = \text{mult}_{\widetilde{h}(\text{univ}_{d_1, \dots, d_t}(\Gamma))}.$$

As a consequence of the surjectivity of the morphism (2.11), we conclude that

$$(4.45) \quad [\widetilde{h}, \text{mult}_x] = \text{mult}_{\widetilde{h}(x)}$$

as an equality of operators $A^*(\text{Hilb}_n) \rightarrow A^*(\text{Hilb}_n)$, indexed by any $x \in A^*(\text{Hilb}_n)$. This is precisely equivalent to (1.6). In the language of correspondences, relation (4.44) is viewed as an equality of correspondences in $A^*(\text{Hilb}_n \times \text{Hilb}_n \times S^t)$. By (2.10) and (2.11), there exist correspondences

$$Z \in A^* \left(\text{Hilb}_n \times \bigsqcup_a S^t \right) \quad \text{and} \quad W \in A^* \left(\bigsqcup_a S^t \times \text{Hilb}_n \right)$$

(here the indexing set follows the notation at the end of Section 2.2) such that $W \circ Z = \Delta_{\text{Hilb}_n}$. By composing the third factor of (4.44) with ${}^t Z \in A^*(\sqcup_a S^t \times \text{Hilb}_n)$, we obtain (4.45) as an equality of correspondences in $A^*(\text{Hilb}_n \times \text{Hilb}_n \times \text{Hilb}_n)$, which is what our result claims.

Let us now prove part (ii), which requires us to show that $\tilde{h}(x) = \deg(x) \cdot x$ if x is a divisor class or a Chern class of the tangent bundle. According to Section 2.5 it suffices to show this relation for

$$x = \text{univ}_{d_1, \dots, d_t}(\Gamma),$$

where $t = 1$ and $\Gamma = \gamma \in \{1, l, c\}_{l \in A^1(S)}$, or $t = 2$ and $\Gamma = \Delta_*(\gamma) \in A^*(S^2)$ for $\gamma \in \{1, c\}$. The degree of such a class x is

$$\deg x = d_1 + \dots + d_t + \deg \Gamma - 2t.$$

As a consequence of (4.43), we have

$$\tilde{h}(x) = \text{univ}_{d_1, \dots, d_t} \left((d_1 + \dots + d_t - t)\Gamma + \int_{\bullet} \sum_{i=1}^t \Gamma_{1\dots i-1, \bullet, i+1\dots t}(c_i - c_{\bullet}) \right)$$

so the class x lies in the appropriate direct summand if

$$\int_{\bullet} \sum_{i=1}^t \Gamma_{1\dots i-1, \bullet, i+1\dots t}(c_i - c_{\bullet}) = (\deg \Gamma - t)\Gamma.$$

If $t = 1$ and $\Gamma = \gamma \in \{1, l, c\}_{l \in A^1(S)}$, this relation is trivial, while if $t = 2$ and $\Gamma = \Delta_*(\gamma)$ for $\gamma \in \{1, c\}$, it is an immediate consequence of Claim 4.5. \square

Proof of Theorem 1.6. The proof follows that of Theorem 1.4 very closely. For part (i) we iterate relations (4.17) and (4.18) to obtain

$$\begin{aligned} [h_{\alpha\beta}, \text{mult}_{\text{univ}_{d_1, \dots, d_t}(\Gamma)}] &= \text{mult}_{\text{univ}_{d_1, \dots, d_t}(\Gamma')}, \\ [h_{\alpha\delta}, \text{mult}_{\text{univ}_{d_1, \dots, d_t}(\Gamma)}] &= \text{mult}_{\sum_{d'_1, \dots, d'_t} \text{univ}_{d'_1, \dots, d'_t}(\Gamma'')}, \end{aligned}$$

where in the right-hand sides, Γ' and Γ'' are obtained from $\Gamma \in A^*(S^t)$ by pulling back to some $S^{t+t'}$, multiplying with certain cycles, and pushing forward to S^t again. In either case, we may apply the relations above to the fundamental class (and invoke Lemma 4.12) to conclude that

$$\begin{aligned} h_{\alpha\beta}(\text{univ}_{d_1, \dots, d_t}(\Gamma)) &= \text{univ}_{d_1, \dots, d_t}(\Gamma'), \\ h_{\alpha\delta}(\text{univ}_{d_1, \dots, d_t}(\Gamma)) &= \sum_{d'_1, \dots, d'_t} \text{univ}_{d'_1, \dots, d'_t}(\Gamma''). \end{aligned}$$

Therefore, we conclude that

$$\begin{aligned} [h_{\alpha\beta}, \text{mult}_{\text{univ}_{d_1, \dots, d_t}(\Gamma)}] &= \text{mult}_{h_{\alpha\beta}(\text{univ}_{d_1, \dots, d_t}(\Gamma))}, \\ [h_{\alpha\delta}, \text{mult}_{\text{univ}_{d_1, \dots, d_t}(\Gamma)}] &= \text{mult}_{h_{\alpha\delta}(\text{univ}_{d_1, \dots, d_t}(\Gamma))}. \end{aligned}$$

As explained at the end of the proof of Theorem 1.4 (i), the formulas above imply (1.10) as an equality of correspondences in $A^*(\text{Hilb}_n \times \text{Hilb}_n \times \text{Hilb}_n)$.

For part (ii), one can check the identity $h_{\alpha\beta}(c_k(\text{Tan}_{\text{Hilb}_n})) = 0$ in the same way as the analogous proof in Theorem 1.4 (ii). Hence it remains to prove that $h_{\alpha\delta}(\text{ch}_k(\text{Tan}_{\text{Hilb}_n})) = 0$. By Lemma 4.12, it suffices to show that the commutator of $h_{\alpha\delta}$ with the operator of multiplication

by $\text{ch}_k(\text{Tan}_{\text{Hilb}_n})$ vanishes. By Section 2.5 this operator is precisely

$$(4.46) \quad \text{mult}_{\text{ch}_k(\text{Tan}_{\text{Hilb}_n})} = 2\mathcal{G}_{k+2}(1) + 4\mathcal{G}_k(c) + \sum_{i+j=k+2} (-1)^{j+1} \mathcal{G}_i \mathcal{G}_j(\Delta) \\ + 2 \sum_{i+j=k} (-1)^{j+1} \mathcal{G}_i(c) \mathcal{G}_j(c).$$

Using (4.15), one has for all d the commutation relations

$$[h_{\alpha\delta}, \mathcal{G}_d(1)] = -\mathcal{G}_2(\alpha) \mathcal{G}_{d-1}(1) - \mathcal{G}_2(1) \mathcal{G}_{d-1}(\alpha) + 2\mathcal{G}_{d-1}(\alpha), \\ [h_{\alpha\delta}, \mathcal{G}_d(c)] = -\mathcal{G}_2(\alpha) \mathcal{G}_{d-1}(c) - \mathcal{G}_{d+1}(\alpha)$$

and for all $i, j \geq 2$ the relations

$$[h_{\alpha\delta}, \mathcal{G}_i \mathcal{G}_j(\Delta)] = -\mathcal{G}_2(\alpha) (\mathcal{G}_{i-1} \mathcal{G}_j + \mathcal{G}_i \mathcal{G}_{j-1})(\Delta) \\ - \mathcal{G}_2(1) (\mathcal{G}_{i-1} \mathcal{G}_j + \mathcal{G}_i \mathcal{G}_{j-1})(\Delta \alpha_1) \\ - (\mathcal{G}_{i+1} \mathcal{G}_j + \mathcal{G}_i \mathcal{G}_{j+1})(\alpha_1 + \alpha_2) \\ + 2\mathcal{G}_{i-1}(\alpha) \mathcal{G}_j(c) + 2\mathcal{G}_i(c) \mathcal{G}_{j-1}(\alpha).$$

By using the above identities, it is straightforward to show that $h_{\alpha\delta}$ commutes with the right-hand side of (4.46) (we leave the computation as an exercise to the interested reader). This implies the required equation, namely $[h_{\alpha\delta}, \text{mult}_{\text{ch}_k(\text{Tan}_{\text{Hilb}_n})}] = 0$. \square

Acknowledgement. We would like to thank Lie Fu, Alina Marian, Davesh Maulik, Junliang Shen, Catharina Stroppel, and Zhiyu Tian for useful discussions.

References

- [1] A. Beauville, Sur l'anneau de Chow d'une variété abélienne, *Math. Ann.* **273** (1986), no. 4, 647–651.
- [2] A. Beauville, On the splitting of the Bloch–Beilinson filtration, in: *Algebraic cycles and motives*. Vol. 2, London Math. Soc. Lecture Note Ser. **344**, Cambridge University, Cambridge (2007), 38–53.
- [3] A. Beauville and C. Voisin, On the Chow ring of a K3 surface, *J. Algebraic Geom.* **13** (2004), no. 3, 417–426.
- [4] M. A. A. de Cataldo and L. Migliorini, The Chow groups and the motive of the Hilbert scheme of points on a surface, *J. Algebra* **251** (2002), no. 2, 824–848.
- [5] C. Deninger and J. Murre, Motivic decomposition of abelian schemes and the Fourier transform, *J. reine angew. Math.* **422** (1991), 201–219.
- [6] L. Fu, R. Laterveer and C. Vial, Multiplicative Chow–Künneth decompositions and varieties of cohomological K3 type, preprint 2019, <http://arxiv.org/abs/1911.06580>; to appear in *Ann. Mat. Pura Appl.*
- [7] L. Fu, R. Laterveer and C. Vial, The generalized Franchetta conjecture for some hyper-Kähler varieties, *J. Math. Pures Appl.* (9) **130** (2019), 1–35.
- [8] L. Fu and Z. Tian, Motivic hyper-Kähler resolution conjecture: II. Hilbert schemes of K3 surfaces, preprint 2017, <http://math.univ-lyon1.fr/~fu/articles/MotivicCrepanthilbK3.pdf>.
- [9] L. Fu and Z. Tian, Motivic multiplicative McKay correspondence for surfaces, *Manuscripta Math.* **158** (2019), no. 3–4, 295–316.
- [10] L. Fu, Z. Tian and C. Vial, Motivic hyper-Kähler resolution conjecture. I: Generalized Kummer varieties, *Geom. Topol.* **23** (2019), no. 1, 427–492.
- [11] L. Fu and C. Vial, Distinguished cycles on varieties with motive of Abelian type and the section property, *J. Algebraic Geom.* **29** (2020), no. 1, 53–107.
- [12] L. Fu and C. Vial, A motivic global Torelli theorem for isogenous K3 surfaces, *Adv. Math.* **383** (2021), Article ID 107674.
- [13] A. Gholampour and R. P. Thomas, Degeneracy loci, virtual cycles and nested Hilbert schemes, I, *Tunis. J. Math.* **2** (2020), no. 3, 633–665.
- [14] M. Green, Y.-J. Kim, R. Laza and C. Robles, The LLV decomposition of hyper-Kähler cohomology, preprint 2019, <http://arxiv.org/abs/1906.03432>.

- [15] *I. Grojnowski*, Instantons and affine algebras. I. The Hilbert scheme and vertex operators, *Math. Res. Lett.* **3** (1996), no. 2, 275–291.
- [16] *K. Künnemann*, A Lefschetz decomposition for Chow motives of abelian schemes, *Invent. Math.* **113** (1993), no. 1, 85–102.
- [17] *R. Laterveer* and *C. Vial*, On the Chow ring of Cynk–Hulek Calabi–Yau varieties and Schreieder varieties, *Canad. J. Math.* **72** (2020), no. 2, 505–536.
- [18] *M. Lehn*, Chern classes of tautological sheaves on Hilbert schemes of points on surfaces, *Invent. Math.* **136** (1999), no. 1, 157–207.
- [19] *W.-P. Li*, *Z. Qin* and *W. Wang*, Hilbert schemes and \mathcal{W} algebras, *Int. Math. Res. Not. IMRN* **2002** (2002), no. 27, 1427–1456.
- [20] *E. Looijenga* and *V.A. Lunts*, A Lie algebra attached to a projective variety, *Invent. Math.* **129** (1997), no. 2, 361–412.
- [21] *E. Markman*, Generators of the cohomology ring of moduli spaces of sheaves on symplectic surfaces, *J. reine angew. Math.* **544** (2002), 61–82.
- [22] *E. Markman*, On the monodromy of moduli spaces of sheaves on $K3$ surfaces, *J. Algebraic Geom.* **17** (2008), no. 1, 29–99.
- [23] *D. Maulik* and *A. Neguș*, Lehn’s formula in Chow and conjectures of Beauville and Voisin, preprint 2019, <http://arxiv.org/abs/1904.05262>; to appear in *J. Inst. Math. Jussieu*.
- [24] *B. Moonen*, On the Chow motive of an abelian scheme with non-trivial endomorphisms, *J. reine angew. Math.* **711** (2016), 75–109.
- [25] *J.P. Murre*, *J. Nagel* and *C.A.M. Peters*, Lectures on the theory of pure motives, *Univ. Lecture Ser.* **61**, American Mathematical Society, Providence 2013.
- [26] *H. Nakajima*, Heisenberg algebra and Hilbert schemes of points on projective surfaces, *Ann. of Math. (2)* **145** (1997), no. 2, 379–388.
- [27] *H. Nakajima*, Lectures on Hilbert schemes of points on surfaces, *Univ. Lecture Ser.* **18**, American Mathematical Society, Providence 1999.
- [28] *A. Neguș*, The Chow of $S^{[n]}$ and the universal subscheme, preprint 2019, <http://arxiv.org/abs/1912.03287>; to appear in *Bull. Math. Soc. Sci. Math. Roumanie*.
- [29] *G. Oberdieck*, A Lie algebra action on the Chow ring of the Hilbert scheme of points of a $K3$ surface, *Comment. Math. Helv.* **96** (2021), no. 1, 65–77.
- [30] *U. Rieß*, On the Chow ring of birational irreducible symplectic varieties, *Manuscripta Math.* **145** (2014), no. 3–4, 473–501.
- [31] *Y. Ruan*, The cohomology ring of crepant resolutions of orbifolds, in: *Gromov–Witten theory of spin curves and orbifolds*, *Contemp. Math.* **403**, American Mathematical Society, Providence (2006), 117–126.
- [32] *M. Shen* and *C. Vial*, The Fourier transform for certain hyperkähler fourfolds, *Mem. Amer. Math. Soc.* **240** (2016), no. 1139, 1–163.
- [33] *M. Shen* and *C. Vial*, The motive of the Hilbert cube $X^{[3]}$, *Forum Math. Sigma* **4** (2016), Paper No. e30.
- [34] *M. Verbitsky*, Cohomology of compact hyper-Kähler manifolds and its applications, *Geom. Funct. Anal.* **6** (1996), no. 4, 601–611.
- [35] *C. Vial*, On the motive of some hyperKähler varieties, *J. reine angew. Math.* **725** (2017), 235–247.
- [36] *C. Voisin*, On the Chow ring of certain algebraic hyper-Kähler manifolds, *Pure Appl. Math. Q.* **4** (2008), no. 3, 613–649.
- [37] *C. Voisin*, Some new results on modified diagonals, *Geom. Topol.* **19** (2015), no. 6, 3307–3343.

Andrei Neguș, Department of Mathematics, MIT, Cambridge, MA, USA;
and Simion Stoilow Institute of Mathematics, Bucharest, Romania
<https://orcid.org/0000-0002-1622-8638>
e-mail: andrei.negut@gmail.com

Georg Oberdieck, Institut für Mathematik, University of Bonn, Bonn, Germany
e-mail: georgo@math.uni-bonn.de

Qizheng Yin, Peking University, BICMR, Beijing, P.R. China
e-mail: qizheng@math.pku.edu.cn

Eingegangen 7. Januar 2020, in revidierter Fassung 28. April 2020